

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
25 November 2004 (25.11.2004)

PCT

(10) International Publication Number
WO 2004/101798 A2

(51) International Patent Classification⁷: C12N 15/82,
C07K 14/01

(21) International Application Number:
PCT/IT2004/000287

(22) International Filing Date: 19 May 2004 (19.05.2004)

(25) Filing Language: Italian

(26) Publication Language: English

(30) Priority Data:
RM2003A000242 19 May 2003 (19.05.2003) IT

(71) Applicants (for all designated States except US):
ENEA-ENTE PER LE NUOVE TECNOLOGIE,
L'ENERGIA E L'AMBIENTE [IT/IT]; Lungotevere
Grande Ammiraglio Thaon di Revel, 76, I-00196 Roma
(IT). CONSIGLIO NAZIONALE DELLE RICERCHE
[IT/IT]; Piazzale Aldo Moro, 7, I-00185 Roma (IT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): TAVAZZA, Mario
[IT/IT]; ENEA-Ente per le nuove Tecnologie, L'Ener-
gia e l'A, mbiente, Lungotevere G.A. Thaon Di Revel,
76, I-00196 Roma (IT). TAVAZZA, Raffaella [IT/IT];
ENEA-Ente per le nuove Tecnologie, L'Energia e l'A,
mbiente, Lungotevere G.A. Thaon Di Revel, 76, I-00196
Roma (IT). LUCIOLI, Alessandra [IT/IT]; ENEA-Ente
per le nuove Tecnologie, L'Energia e l'A, mbiente, Lun-
gotevere G.A. Thaon Di Revel, 76, I-00196 Roma (IT).
BRUNETTI, Angela [IT/IT]; ENEA-Ente per le nuove
Tecnologie, L'Energia e l'A, mbiente, Lungotevere G.A.
Thaon Di Revel, 76, I-00196 Roma (IT). BERARDI,
Alessandra [IT/IT]; ENEA-Ente per le nuove Tecnolo-
gie, L'Energia e l'A, mbiente, Lungotevere G.A. Thaon

Di Revel, 76, I-00196 Roma (IT). NORIS, Emanuela
[IT/IT]; Consiglio Nazionale delle Ricerche, Pl.le Aldo
Moro, 7, I-00185 Roma (IT). ACCOTTO, Gian, Paolo
[IT/IT]; Consiglio Nazionale delle Ricerche, Pl.le Aldo
Moro, 7, I-00185 Roma (IT).

(74) Agents: GITTO, Serena et al.; Ing. Barzanò & Zanardo
Roma S.p.A., Via Piemonte, 26, I-00187 Roma (IT).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR THE PREPARATION OF TRANSGENIC PLANTS CHARACTERISED BY GEMINIVIRUS LAST-
ING RESISTANCE

(57) Abstract: The invention relates to a method for the production of transgenic plants durably resistant against *Geminiviruses*,
wherein the transgene consists of a suitably modified polynucleotide sequence, derived by the pathogen.

WO 2004/101798 A2

DESCRIPTION

METHOD FOR THE PREPARATION OF TRANSGENIC PLANTS
CHARACTERISED BY GEMINIVIRUS LASTING RESISTANCE

5

* * *

The present invention relates to a method for the preparation of transgenic plants lasting resistant to geminiviruses.

10 More particularly the invention concerns a method for the preparation of transgenic plants lasting resistant to geminiviruses, wherein the transgene consists of a polynucleotide sequence, derived from the pathogen, suitably modified in order to result in an ineffective target of the post-transcriptional gene silencing induced by geminiviruses.

15 It is known that geminiviruses are a wide and diversified class of plant viruses that infect several plants of agronomic interest causing serious harvest losses. Such viruses are characterised by virions consisting of two geminate icosahedric particles. Their genome, consisting of one or two circular single-stranded DNA molecules (ssDNA), replicates in the nucleus of infected cells through double stranded intermediates (Hanley-Bowdoin et al., 1999).

20 The *Geminiviridae* family is divided in four genera named *Mastrevirus*, *Begomovirus*, *Curtovirus* and *Topocuvirus* based on the insect vector, the host spectrum and the genome structure (Briddon et al., 1985; Fauquet et al., 2003).

25 A serious disease of the tomato plant, transmitted by the whitefly *Bemisia tabaci*, is from a long time known as "tomato yellow leaf curl" in the areas of the Middle East, Asian South East and Africa, (Czosnek et al., 1997). This disease, that can cause harvest losses of 100% (Picò et al., 1996; Czosnek et al., 1997), successively spread both throughout the Western Mediterranean, reaching Sardinia, Sicily and Spain (Czosnek et al., 1997), and America (Polston et al., 1997).

30 Recently the agents of the disease have been identified and isolated, being viruses belonging to the *Geminiviridae* family, genera *Begomovirus*. Phylogenetic studies have highlighted the presence of different viral species related to different geographical origins of the *Begomovirus*: Asia, Africa and America (Czosnek et al., 1997).

The genome of the Tomato yellow leaf curl Sardinia virus (TYLCSV) species, is monopartite (Kheyr-Pour et al., 1991). The DNA is transcribed

bidirectionally and contains six open reading frame (ORF), two on the viral strand (V): V1 and V2, and four on the complementary strand (C): C1, C2, C3 and C4, as shown in figure 1. Between the C1 and V2 ORFs there is a non-coding region named intergenic region (IR) analogous to that present in the genome of all *Geminiviridae*. The genomic organization of TYLCSV is structurally similar to that of the bipartite *Begomoviruses* component A such as the tomato golden mosaic virus (TGMV) and the African cassava virus (ACMV). In the case of bipartite *Begomoviruses* the nomenclature of the ORFs present on the component A of the complementary strand is: AL1 or AC1, AL2 or AC2, AL3 or AC3, AL4 or AC4, while on the viral strand AR1 or AV1, AR2 or AV2; on the complementary strand of the component B is: BL1 or BC1 and on the viral strand BR1 or BV1.

Strategies used until now in order to control the infection of the geminiviruses transmitted by the *Bemisia tabaci* are based on the use of expensive fine mesh nets (for the cultivation of fresh-market tomato) and particularly on repeated insecticide treatments (cultivations of both fresh-market and processing tomato). Such strategies result in an increase of the production expenses and represent a serious danger for the health of the agricultural operators and consumer. Furthermore the onset of *Bemisia tabaci* populations resistant to the insecticide imidacloprid has been already reported (Cahill et al., 1996; Williams et al., 1996).

The development of resistant cultivated species represents the most practical and economic way to control viral infections. Classical breeding programs for introducing resistance to geminiviruses that cause the tomato yellow leaf curl were based on the transfer of resistance genes from wild species of *Lycopersicon* to species of cultivated tomato. Thereby lines with variable levels of resistance to TYLCSV have been obtained and commercialized, the best lines showing reduced symptoms and low viral replication. However plants with low and mean levels of resistance represent a potential receptacle for further infections.

Another important aspect to be considered is that the agronomic characteristics of the lines obtained are not always optimal and however reflects those of the genotype of cultivated tomato used in breeding programs.

A tomato line immune to the viruses causing the tomato yellow leaf curl disease, namely, with neither symptoms nor viral DNA replication has not been released yet.

With the advent of genetic engineering new perspectives were opened up for the introduction of resistance characters against plant viruses. Most strategies are based on the introduction and expression of pathogen-derived sequences in the plant of interest, Pathogen Derived Resistance (PDR) (Sanford & Johnson, 1985; Abel et al., 1986; Tavazza and Lucioli, 1993).

Although such strategies have been successfully applied for the introduction of resistance characters to plant viruses with RNA genome (Beachy, 1997), in the case of geminiviruses, with a DNA genome, the expression of pathogen-derived sequences has produced plants with no lasting resistance and/or tolerance.

The mechanisms that induce virus resistance achieved through the expression of pathogen-derived sequences can be grouped in two wide classes:

- a) resistance mediated by the expression of a pathogen protein such as, for instance, the expression of a dominant negative mutant;
- b) resistance mediated by the post-transcriptional gene silencing (Baulcombe, 1996; Beachy, 1997; Zaitlin and Palukaitis, 2000).

The post-transcriptional gene silencing is a ubiquitary process in eukaryotes, involving the degradation of specific RNAs following the formation of double strand RNA (dsRNA) molecules having sequences homologous to the target RNA.

Although there may be different contexts able to induce the production of dsRNA homologous to the transgene (transcription of aberrant transgenic RNAs, presence in the transgenic RNA of sufficiently long inverted and repeated sequences, integration of the transgene in the plant genome in inverted and repeated multiple copies), once the dsRNA is produced, the latter is recognised and degraded in short molecules of dsRNA of about 21-26 nucleotides, referred to as siRNA.

The siRNAs are then integrated in a multiprotein complex named RISC, that is able to degrade all RNAs having sequence homology with the siRNAs. The latter ones represent therefore the determining factors of RNA silencing specificity and their presence related to a determined sequence establishes univocally that this RNA sequence is post-transcriptionally silenced.

Therefore, transgenic plants post-transcriptionally silenced for sequences derived from viral RNA genome, are resistant to the

homologous virus and to viruses with nucleotide sequences closely related to the transgene.

The transgene silencing can be also induced following virus infection.

5 In fact, viral replication is able to induce silencing of a transgene, initially not silenced, if the nucleotide sequence of the transgene is homologous to a portion of the infecting virus genome. The activation of the silencing mechanism involves the specific degradation of the RNA molecules having sequence homology with the inducer RNA.

10 As direct consequence, the silencing activation by the virus is associated with a degradation of both transgenic mRNA sequences homologous to the virus and viral genome. This results in the host recovery after an initial infectious step, so that the new vegetative part is proved to be virus free. A peculiar characteristic of the plant tissues that
15 develop subsequently to the recovery phenomenon is that they are highly resistant to a following infection by the same virus.

The resistance mediated by post-transcriptional gene silencing, since based on recognition at the nucleotidic level, confers resistance only against viral isolates closely homologous to the virus genome from
20 which the transgene was derived. Instead, strategies based on the expression of a pathogen protein normally produce plants resistant also to viral strains or isolates not-closely related from a nucleotide point of view.

It is also been shown that the transgene silencing is influenced by the temperature, being inactive at temperatures below 15°C (Szittyá et al., 2003). Therefore plants exposed in field conditions at temperature
25 range below 15°C can lose the silencing-mediated resistance.

It must be borne in mind that, although from several years transgenic plants resistant to RNA genome viruses have been achieved through mechanisms based on transgene silencing, so far it is not
30 reported that such strategy can be successfully applied to the geminiviruses (DNA genome-viruses).

It's clear that the best strategy in order to obtain plants resistant to a wide spectrum of geminiviruses is the one in which the interfering product is the protein. It is clear that the width of the resistance spectrum
35 increases the agronomic and commercial value of the produced plant.

Thereby the expression in transgenic plants of dysfunctional variants of geminivirus replicative Rep protein has been used in order to

obtain plants with greater levels of resistance or immunity against the geminiviruses.

It's known in literature that the expression of a truncated replicative Rep protein (Rep-210) of TYLCSV is able to confer resistance against viral infection, although such resistance is not lasting because the virus is able to overcome it over time.

In tables 1 and 2 are shown the results of the analysis of the resistance of TYLCSV-agroinoculated Rep-210 expressing transgenic plants of *Tomato* 47 x wt (Brunetti et al. 1997) and of *N. benthamiana* line 102.22 (Noris et al. 1996) respectively.

Table 1

<i>Lycopersicon esculentum</i>	Time (weeks)	N° infected plants/inoculated plants	% infected plants/inoculated plants
Rep-210	4	0/13	0
	9	2/13	15
	18	5/13	38
Wild-type	4	6/6	100

Table 2

<i>Nicotiana benthamiana</i>	Time (weeks)	N° infected plants/inoculated plants	% infected plants/inoculated plants
Rep-210	2	4/21	19
	3	11/21	52
	4	18/21	86
Wild-type	2	6/6	100

From the results reported in tables 1 and 2, it can be clearly inferred that the resistance against TYLCSV mediated by the transgenic expression of a pathogen-derived sequence, is overcome with time.

Similarly, also the resistance induced by the transgenic expression of a dominant negative mutant of Rep of the bipartite geminivirus "African Cassava Mosaic Virus" is overcome with time (Sangaré et al., 1999).

Another example is represented by the transgenic expression of the TYLCV capsid protein in a tomato interspecific hybrid (*Lycopersicon esculentum* X *L. pennellii*) which confers a partial resistance against viral infection (Kunik et al., 1994). Even in this case the resistance mediated by

the expression of the capsid protein is not long lasting and it results to be poorly useful from an agronomic point of view.

In the light of the above, is clear the need to have new methods that would allow to use successfully the polynucleotide sequences derived from the geminiviruses in order to obtain long lasting resistant plants against geminiviruses.

The authors of the present invention have now prepared polynucleotide sequences encoding pathogen-derived viral proteins and able to confer virus resistance to the host, suitably modified in order to be ineffective targets of the virus-induced post-transcriptional gene silencing to obtain transgenic plants with lasting levels of resistance against geminiviruses.

In fact during the experiments the authors show that the overcoming of the resistance, and therefore the difficulty to achieve lasting resistance against geminiviruses, is due to the unexpected abilities of the geminiviruses to silence post-transcriptionally the transgene and to spread in a plant in which the transgene, with sequences homologous to the infecting virus, is post-transcriptionally silenced.

As shown in figures 2 and 3, respectively, both in the transgenic plants of *N. benthamiana* line 102.22 and in the plants of *Tomato 47 x wt* the virus ability to overcome the resistance results from the transgene silencing by the same virus and from the unexpected ability of the virus to spread in a silenced plant.

The TYLCSV ability to spread in a plant in which the transgene Rep-210 is post-transcriptionally silenced, is further circumstantiated as set forth in figure 4.

The results show that the transgenic *tomato* plants *47 x 10D* (Brunetti et al., 1997), post-transcriptionally silenced before agroinoculation, as shown by the absence of the Rep-210 protein and by the concurrent presence of the transgene-homologous siRNAs, are susceptible to the TYLCSV infection as well as the controls.

From the above it results that, contrary to RNA viruses, the geminivirus is not blocked by an active silencing of viral gene sequences. The above said is not limited to the kind of transgenic plant to be used or the way the virus should be inoculated, through agroinfection or *Bemisia tabaci*. In fact, as shown in table 3, using a reduced number of viruliferous *bemisia* per plant, so as to infect between 90% and 100% of the control

plants, about 40% of transgenic plants (line 201) whose transgene is post-transcriptionally silenced, are not or late infected, while at a higher inoculum concentration, all the plants challenged with viruliferous insects are infected similarly to the experiments carried out using agroinoculation.

5

Table 3

Molecular Analysis before inoculum		Low concentration of inoculum ^a			High concentration of inoculum ^b		
Transgenic plants		2 ^c	3	6	2	3	6
Rep-210 (No)	siRNAs (Si)	6/15	7/15	8/15	16/21	20/21	21/21
Not transgenic		2 ^c	3	6	2	3	6
Rep-210 (No)	siRNAs (No)	11/12	11/12	11/12	8/8	8/8	8/8

a Seven viruliferous insects per plant for 2 days

b Thirty-five viruliferous insects per plant for 5 days

c Weeks after inoculum

Therefore it's important to consider that the viral agroinoculation conditions used for testing the resistance and assessing persistence over time (as shown in figures 2, 3 and 4 and in tables 1 and 2) correspond to high or very high viral pressure conditions. This experimental approach allows to identify transgenic plants with very high resistance levels or immune against the viral infection and therefore of very high commercial value.

15

Accordingly, the introduction of resistance characters against geminiviruses through the expression of pathogen-derived sequences is limited due to the unexpected ability of the geminiviruses to silence post-transcriptionally the transgene and to spread in the silenced plant.

20

Furthermore the authors show that the transcripts both of positive (V1 and V2) and negative strand (C1, C2, C3 and C4) of TYLCSV are subjected, during a normal infection on wild-type plants, to the viral post-transcriptional silencing, as shown in figure 5. This results in the impossibility to achieve long-term resistance through expression of sequences derived from the same pathogen, unless these are suitably modified in order not to be a target or to be an ineffective target of the virus-induced post-transcriptional gene silencing. Instead, by introducing in the plant genome a sequence suitably mutated or chosen according to the

25

invention it is possible to obtain a long lasting resistance against geminiviruses, unlike that achieved with the known methods.

Therefore it is an object of the present invention a polynucleotide sequence encoding an aminoacidic sequence derived from
5 geminiviruses, said polynucleotide sequence being characterised in that it is not a target or it is an ineffective target of the viral post-transcriptional silencing and having:

a) a nucleotide homology lower or equal to 90% with respect to the corresponding gene sequence of the geminiviruses against which a
10 resistance is required, preferably lower or equal to 80%, more preferably lower or equal to 70 %;

b) a continuous homology in the RNA transcript, with respect to the corresponding gene sequence of the geminiviruses against which a resistance is required, lower or equal to 17 nucleotides, preferably lower or
15 equal to 8 nucleotides, more preferably lower or equal to 5 nucleotides;

c) a maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of the geminiviruses no longer than 30 nucleotides, preferably no longer than 20 nucleotides, more preferably equal or lower than 9 nucleotides;

20 said polynucleotide sequence being able to confer to the whole plants, tissues or plant cells therewith transformed, a lasting resistance against the geminiviruses.

The polynucleotide sequences according to the invention can be wild-type or synthetic or produced by mutagenesis and the geminivirus-derived aminoacidic sequences encoded by them are wild-type or mutant
25 sequences that interfere with the viral infection.

Therefore the invention includes polynucleotide sequences of geminivirus either suitably changed or wild-type, such as to differ, at the nucleotidic level, with respect to the corresponding genomic sequence of
30 the geminivirus against which it is required to introduce resistance according to the principles above defined and specified in a), b) and c).

Further object of the present invention is a polynucleotide sequence encoding a geminivirus-derived aminoacidic sequence, said polynucleotide sequence being characterised in that it is not a target or it
35 is an ineffective target of the post-transcriptional silencing and having homology even equal to 100% with respect to the sequence of the geminivirus against which it is required a resistance and being shortened

so as to be underrepresented in the siRNAs population with respect to the original sequence, even if maintaining similar interfering abilities.

The gene sequences from which constructing the polynucleotide sequence according to the invention can derive from the geminiviruses such as, *Mastrevirus*, *Curtovirus*, *Begomovirus*, *Topocuvirus* and particularly can be derived from the species shown in table 4 and their isolates, more particularly from the species of Tomato yellow leaf curl and their isolates shown in table 5.

Table 4

List of geminivirus species	Acronym
African cassava mosaic virus	ACMV
Bean calico mosaic virus	BcaMV
Bean dwarf mosaic virus	BDMV
Bean golden mosaic virus	BGMV
Bean golden yellow mosaic virus	BGYMV
Cabbage leaf curl virus	CaLCuV
Chilli leaf curl virus	ChiLCuV
Cotton leaf crumple virus	CLCrV
Cotton leaf curl Alabad virus	CLCuAV
Cotton leaf curl Gezira virus	CLCuGV
Cotton leaf curl Kokhran virus	CLCuKV
Cotton leaf curl Multan virus	CLCuMV
Cotton leaf curl Rajasthan virus	CLCuRV
Cowpea golden mosaic virus	CPGMV
Cucurbit leaf curl virus	CuLCuV
East African cassava mosaic Cameroon virus	EACMCV
East African cassava mosaic Malawi virus	EACMMV
East African cassava mosaic virus	EACMV
East African cassava mosaic Zanzibar virus	EACMZV
Indian cassava mosaic virus	ICMV
Ipomea yellow vein virus	IYVV
Melon chlorotic leaf curl virus	MCLCuV
Mungbean yellow mosaic India virus	MYMIV
Mungbean yellow mosaic virus	MYMV
Okra yellow vein mosaic virus	OYVMV
Papaya leaf curl virus	PaLCuV
Pepper golden mosaic virus	PepGMV

Pepper huasteco yellow vein virus	PHYVV
Pepper leaf curl Bangladesh virus	PepLCBV
Pepper leaf curl virus	PepLCV
Potato yellow mosaic Panama virus	PYMPV
Potato yellow mosaic Trinidad virus	PYMTV
Potato yellow mosaic virus	PYMV
South African cassava mosaic virus	SACMV
Soybean crinkle leaf virus	SbCLV
Squash leaf curl China virus	SLCCNV
Squash leaf curl virus	SLCV
Squash leaf curl Yunnan virus	SLCYV
Squash mild leaf curl virus	SMLCV
Squash yellow mild mottle virus	SYMMoV
Sri Lankan cassava mosaic virus	SLCMV
Sweet potato leaf curl Georgia virus	SPLCGV
Sweet potato leaf curl virus	SPLCV
Tobacco curly shoot virus	TbCSV
Tobacco leaf curl Japan virus	TbLCJV
Tobacco leaf curl Kochi virus	TbLCKoV
Tobacco leaf curl Yunnan virus	TbLCYNV
Tobacco leaf curl Zimbabwe virus	TbLCZV
Tomato chlorotic mottle virus	ToCMoV
Tomato golden mosaic virus	TGMV
Tomato golden mottle virus	ToGMoV
Tomato leaf curl Bangalore virus	ToLCBV
Tomato leaf curl Bangladesh virus	ToLCBDV
Tomato leaf curl Gujarat virus	ToLCGV
Tomato leaf curl Karnataka virus	ToLCKV
Tomato leaf curl Laos virus	ToLCLV
Tomato leaf curl Malaysia virus	ToLCMV
Tomato leaf curl New Delhi virus	ToLCNDV
Tomato leaf curl Sri Lanka virus	ToLCSLV
Tomato leaf curl Taiwan virus	ToLCTWV
Tomato leaf curl Vietnam virus	ToLCVV
Tomato leaf curl virus	ToLCV
Tomato mosaic Havana virus	ToMHV
Tomato mottle Taino virus	ToMoTV

Tomato mottle virus	ToMoV
Tomato rugose mosaic virus	ToRMV
Tomato severe leaf curl virus	ToSLCV
Tomato severe rugose virus	ToSRV
Tomato yellow leaf curl China virus	TYLCCNV
Tomato yellow leaf curl Gezira virus	TYLCGV
Tomato yellow leaf curl Malaga virus	TYLCMaIV
Tomato yellow leaf curl Sardinia virus	TYLCSV
Tomato yellow leaf curl Thailand virus	TYLCTHV
Tomato yellow leaf curl virus	TYLCV
Watermelon chlorotic stunt virus	WmCSV
Wheat dwarf virus	WDV
Maize streak virus	MSV
Sugarcane streak virus	SSV
Bean yellow dwarf virus	BYDV
Tobacco yellow dwarf virus	TYDV
Tomato pseudo curly top virus	TPCTV
Beet curly top virus	BCTV

Table 5

Species of tomato yellow leaf curl (Fauquet <i>et al.</i> , 2003)	Acronym
<i>Tomato yellow leaf curl China virus</i>	TYLCCNV
Tomato yellow leaf curl China virus AF311734	TYLCCNV
Tomato yellow leaf curl China virus – [Y64] AJ457823	TYLCCNV-[Y64]
Tomato yellow leaf curl China virus – Tb [Y10] AJ319675	TYLCCNV-Tb[Y10]
Tomato yellow leaf curl China virus – Tb [Y11] AJ319676	TYLCCNV-Tb[Y11]
Tomato yellow leaf curl China virus – To [Y25] AJ457985	TYLCCNV-Tb[Y25]
Tomato yellow leaf curl China virus – Tb [Y36] AJ420316	TYLCCNV-Tb[Y36]
Tomato yellow leaf curl China virus – Tb [Y38] AJ420317	TYLCCNV-Tb[Y38]
Tomato yellow leaf curl China virus – Tb [Y5] AJ319674	TYLCCNV-Tb[Y5]
Tomato yellow leaf curl China virus – Tb [Y8] AJ319677	TYLCCNV-Tb[Y8]

<i>Tomato yellow leaf curl Gezira virus</i>	TYLCGV
Tomato yellow leaf curl Gezira virus – [1] AY044137	TYLCGV-[1]
Tomato yellow leaf curl Gezira virus – [2] AY044138	TYLCGV-[2]
Tomato yellow leaf curl Gezira virus – [Shambat] AY044139	TYLCGV-[Sha]
<i>Tomato yellow leaf curl Malaga virus</i>	TYLCMaIV
Tomato yellow leaf curl Malaga virus AF271234	TYLCMaIV
<i>Tomato yellow leaf curl Sardinia virus</i>	TYLCSV
Tomato yellow leaf curl Sardinia virus X61153	TYLCSV
Tomato yellow leaf curl Sardinia virus – Spain [1] Z25751	TYLCSV-ES[1]
Tomato yellow leaf curl Sardinia virus – Spain [2] L27708	TYLCSV-ES[2]
Tomato yellow leaf curl Sardinia virus – Sicily Z28390	TYLCSV-Sic
<i>Tomato yellow leaf curl Thailand virus</i>	TYLCTHV
Tomato yellow leaf curl Thailand virus – [1] X63015, X63016	TYLCTHV-[1]
Tomato yellow leaf curl Thailand virus – [2] AF141922, AF141897	TYLCTHV-[2]
Tomato yellow leaf curl Thailand virus – [Myanmar] AF206674	TYLCTHV-[MM]
Tomato yellow leaf curl Thailand virus – [Y72] AJ495812	TYLCTHV-[Y72]
<i>Tomato yellow leaf curl virus</i>	TYLCV
Tomato yellow leaf curl virus X15656	TYLCV
Tomato yellow leaf curl virus – [Almeria] AJ489258	TYLCV-[Alm]
Tomato yellow leaf curl virus – [Aichi] AB014347	TYLCV-[Aic]
Tomato yellow leaf curl virus – [Cuba] AJ223505	TYLCV-[CU]
Tomato yellow leaf curl virus – [Dominican Republic] AF024715	TYLCV-[DO]
Tomato yellow leaf curl virus – [Portugal] AF105975	TYLCV-[PT]
Tomato yellow leaf curl virus – [Saudi Arabia]	TYLCV-[SA]
Tomato yellow leaf curl virus – [Shizuokua] AB014346	TYLCV-[Shi]
Tomato yellow leaf curl virus – [Spain7297] AF071228	TYLCV-[ES7297]
Tomato yellow leaf curl virus – Iran AJ132711	TYLCV-IR
Tomato yellow leaf curl virus – Mild X76319	TYLCV-Mld

5 Preferably the species of *Begomovirus* are TYLCCNV, TYLCGV, TYLCMaIV, TYLCSV, TYLCTHV, TYLCV, ACMV, BGMV, CaLCuV, ToCMoV, TGMV, ToGMOV, ToMHV, ToMoTV, ToMoV, ToRMV, ToSLCV, ToSRV, Cotton leaf curl (CLCrV, CLCuAV, ClCuGV, CLCuKV, CLCuMV, CLCuRV), East African cassava mosaic (EACMCV, EACMMV, EACMV, EACMZV), Potato yellow mosaic (PYMPV, PYMTV, PYMV), Squash leaf curl (SLCCNV, SLCV, SLCYV), Sweet potato leaf curl (SPLCGV, SPLCV), Tobacco leaf curl (TbLCJV, TbLCKoV, TbLCYNV, TbLCZV), Tomato leaf

curl (ToLCBV, ToLCBDV, ToLCGV, ToLCKV, ToLCLV, ToLCMV, ToLCNDV, ToLCSLV, ToLCTWV, ToLCVV, ToLCV) and isolates thereof.

Other species of preferred geminivirus, belonging to the other genera *Mastrevirus*, *Curtovirus*, *Topocuviruses*, are WDV, MSV, SSV, BYDV, TYDV, BCTV and their isolates.

The gene sequence belonging to the genome of the geminiviruses can be the sequence C1/AL1/AC1, C2/AL2/AC2, C3/AL3/AC3, C4/AL4/AC4, V1/AR1/AV1, V2/AR2/AV2, BC1/BL1 and BV1/BR1, particularly, the sequence C1/AL1/AC1 of the previously described geminiviruses and their isolates.

The aminoacidic sequence encoded by the polynucleotide sequence object of the present invention is a pathogen-derived protein able to confer resistance against the geminiviruses to the plants expressing it. Said interfering protein since, according to the invention, is stably expressed, confers a lasting resistance independently from the molecular mechanism by which the protein product is able to induce resistance.

The pathogen-derived protein can be a capsid protein, replication-associated viral protein (Rep), proteins encoded by the genes C2/AL2/AC2, C3/AL3/AC3, C4/AL4/AC4, V2/AR2/AV2, BC1/BL1 and BV1/BR1.

An example of a possible polynucleotide sequence satisfying the above reported requirement is set forth in figures 16A and 16B that show the alignment between the wild-type nucleotide sequence encoding the Rep-210 protein of the TYLCSV and the synthetic nucleotide sequence modified so as not to be a target of the post-transcriptional degradation induced by the infecting virus, where both nucleotide sequences encode the same viral protein.

The plants, tissues or plant cells that can be transformed with this polynucleotide sequences can be tomato, pepper, tobacco, sweet potato, cotton, melon, squash, manioc, potato, bean, soybean, mung bean, beet, sugar cane, corn, wheat.

It is a further object of the present invention a construct comprising an heterologous polynucleotide sequence containing in 5'-3' direction:

a) a polynucleotide sequence acting as promoter in said plant or tissue or transformed cells;

b) a non-translated polynucleotide sequence positioned at 5' of the encoding region, belonging or not to the intergenic region of geminivirus;

5 c) a polynucleotide sequence according to the invention or a fragment or a variant thereof;

d) a sequence acting as terminator of transcription, positioned at the 3' with respect to said polynucleotide sequence.

A further object of the present invention is an expression vector comprising the previously described construct.

10 Further it is an object of the present invention a plant, tissue or transgenic plant cells, progeny thereof as well as seeds comprising in their genome a polynucleotide sequence according to the present invention.

Finally, it is an object of the present invention a method for the preparation of transgenic plants, tissues or plant cells thereof long-lasting
15 resistant to the geminiviruses that comprises the following steps:

a) "identification" or "selection" of a viral gene sequence encoding an aminoacidic sequence able to confer resistance against geminiviruses;

20 b) mutagenesis or "choice" of the viral gene sequence so as to make it an ineffective target of the post-transcriptional silencing induced by infecting geminivirus;

c) insertion of the geminivirus mutated or chosen gene sequence obtained in step b) through a construct as described previously, in the plant, tissue or plant cell thereof.

25 With reference to step a) of the method according to the present invention, the term "identification" means the experimental recognition of said viral gene sequence able to confer resistance against geminiviruses, while the term "selection" means the recognition of an already available viral gene sequence able to confer a not lasting
30 resistance against geminiviruses. Accordingly, the method according to the present invention provides furthermore the solution to the problem of the loss of resistance against geminiviruses that occurs through the employment of known sequences.

35 Particularly, the mutagenesis predicted in step b) is carried out maintaining a nucleotide homology, with respect to the corresponding gene sequence of the geminiviruses against which it is required to obtain a resistance, lower or equal to 90%, preferably lower or equal to 80%, more

preferably lower or equal to 70%, distributed so as the continuous homology in the transcribed RNA with respect to the corresponding sequence of geminiviruses is lower or equal to 17 nucleotides, preferably lower or equal to 8 nucleotides, more preferably lower or equal to 5
5 nucleotides and the maximum length of the sequence containing a single substitution with respect to the native gene sequence is not more than 30 nucleotides, preferably not more than 20 nucleotides, more preferably lower or equal to 9 nucleotides.

As the aminoacidic sequence encoded by the polynucleotide
10 sequence identified or selected in step a), according to the present invention, it can be a protein having homology of 100% with respect to the viral wild-type protein.

This mutagenesis includes all those mutations on the nucleotide sequence that don't decrease the ability of the protein to confer resistance
15 against geminivirus. Possible mutations are both silent point mutations and those leading to the substitution with aminoacids having similar biochemical characteristics, or deletions and/or insertions and/or substitutions.

Alternatively, the mutagenesis in step b) of the method
20 according to the present invention consists of deletions of the polynucleotide sequence at the extremities so as said sequence, while maintaining similar interfering abilities, is under-represented with respect to the original sequence, in the natural population of the siRNAs produced by the infecting virus.

Alternatively the "choice" in step b) of the method according to
25 the present invention consists in the recognition of geminivirus wild-type sequences that differ at the nucleotidic level from the geminivirus against which it is required resistance so as not to be a target or to be an ineffective target of the post-transcriptional silencing.

Particularly, the mutagenesis action in step b) of the method
30 according to the present invention can consist of deletions of the 3' or 5' region of the viral gene sequence of step a), until it is identified the minimum region of said gene sequence that is under-represented with respect to the sequence encoding a wild-type protein, in the population of
35 the siRNAs and that said truncated protein maintains the ability to confer resistance against geminiviruses.

Moreover, the viral gene sequence of step a) of the method according to the present invention can be that of TYLCSV C1/AL1/AC1 gene and the aminoacid sequence can be a protein truncated relatively to the viral wild-type protein such as, for instance, Rep-130.

5 Among various agronomic applications of the synthetic polynucleotide sequences according to the present invention, of particular interest is their use for obtaining tomato plants resistant to TYLCSV.

In this particular embodiment, the transgenic polynucleotide sequence encoding the truncated viral Rep protein (Rep-210) has been
10 modified through a 3' deletion resulting in an ineffective target of TYLCSV-induced post-transcriptional gene silencing, while maintaining the ability to confer resistance.

In particular, using stringent hybridizations with radioactive RNA probes, it was identified a transcribed region of the TYLCSV genome that
15 is under-represented in the population of viral-origin siRNAs produced during the infection of the TYLCSV in wild-type plants, as shown in figures 6' and 7. This region corresponds to the first 390 nucleotides of the gene encoding the TYLCSV Rep. Its transcript is an ineffective target of virus-induced post-transcriptional gene silencing, it encodes the Rep-130 protein
20 which results stably expressed, resulting in a lasting resistance.

Therefore, in a particular embodiment of the invention, the aminoacid sequence of geminiviruses, such as the TYLCSV, encoded by the polynucleotide sequence according to the invention can be the truncated Rep-130 protein (SEQ ID No 9). In this case the viral gene
25 sequence made an ineffective or non target of the post-transcriptional silencing, is the SEQ ID No 8.

It's a further object of the present invention a method as described above wherein the mutagenesis in step b) consists of silent point mutations of the viral gene sequence of step a) that maintain the
30 ability of the encoded aminoacid sequence, to confer resistance against geminiviruses and to be an ineffective or non target of the post-transcriptional silencing.

Particularly, the viral gene sequence of step a) can be the V1/AR1/AV1 (CP) gene for instance of TYLCSV (SEQ ID No 12), and in a
35 particular embodiment the viral gene sequence made ineffective or non target of the post-transcriptional silencing is the SEQ ID No 6.

In addition, the viral gene sequence of step a) can be the TYLCSV C1/AL1/AC1 gene, and in this case the viral gene sequence which was made ineffective or non target of the post-transcriptional silencing can be the SEQ ID No 2 or the SEQ ID No 4.

5 The present invention now will be described by way of illustrating but not limiting way, according to preferred embodiments thereof, with particular reference to the figures of the enclosed drawings, wherein:

10 figure 1 shows the genome of tomato yellow leaf curl Sardinia specie virus (TYLCSV). DNA is transcribed bidirectionally and it contains six open reading frames partially or totally overlapping (ORF), two on the viral strand (V) V1 and V2 and four on the complementary strand (C) C1, C2, C3 and C4;

15 figure 2 shows the expression of Rep-210 protein in TYLCSV-agroinoculated transgenic *N. benthamiana* plants (line 102.22). The symbols (-), (+) and NI mean healthy, infected and non-inoculated plants, respectively. Analysis has been carried out before the agroinoculation with TYLCSV (0 wpi) and four and eight weeks after it (respectively 4 and 8 wpi);

20 figure 3 shows the expression of the Rep-210 protein and of the transgenic mRNAs in tomato plants (line 47 X wt) before the agroinoculation with TYLCSV (0 wpi) and 22 weeks after it (22 wpi). The symbols (+) and (-) mean plants that are, respectively, infected or healthy at the specified time;

25 figure 4 shows the analysis of expression of Rep-210 protein and of the siRNAs corresponding to the relative transcript in transgenic tomato plants (47 X 10D line) before TYLCSV agroinoculation. The symbols (+) and (-) on the panel mean the presence and absence of the sense and antisense C1 transgene, respectively; wt means control wild-type plant; while the symbols (+), (-) and NI under the panel of the siRNAs mean the presence (+) and absence (-), respectively, of the virus and the non-agroinoculation (NI) as a control;

30 figure 5 shows the Northern blot of the small RNAs extracted from TYLCSV-infected wild-type tomato plants (samples 1-4) and non-infected as a control (sample C); M means a molecular weight marker;

figure 6 shows the distribution analysis of the small interfering RNAs with respect to the genome of the TYLCSV. On the top, the linear

map of the TYLCSV genome; transcripts are pointed out by the arrows (V1 and V2 with the same polarity as the viral genome and C1, C2, C3 and C4 with complementary polarity); open boxes from 1 to 9 positioned under the viral genome map represent the nine PCR fragments (each about three hundred nucleotides in length), in which the genome has been divided and of which the ethidium bromide staining and hybridization with the siRNAs extracted by TYLCSV-infected tomato plants are shown. IR represents a tenth PCR fragment, corresponding to the non-transcribed intergenic region of the TYLCSV genome. The numbers under the panels mean the percentage of hybridization signal with respect to each PCR fragment;

figure 7 shows the analysis of the presence of siRNAs in non-agroinoculated (sample C) or TYLCSV-agroinoculated (samples 1 and 2) wild-type tomato plants at four weeks after inoculation; particularly the siRNAs corresponding to the transcript for Rep-210 (probe A) and for Rep-130 (probe B) were analyzed; the columns 100 and 50 on the left panels correspond to 100 and 50 pg, respectively, of an oligonucleotide homologous to both the A and B probes;

figure 8 shows the nucleotide sequence (SEQ ID No 8) encoding Rep-130 (SEQ ID No 9) of the pTOM130 plasmid. In capital, nucleotides not belonging to the TYLCSV but deriving from cloning; the underlined sequences correspond to BamHI and EcoRI restriction sites used for cloning. The start and stop codons are set forth in boldfaced while the mutations introduced for eliminating the C4 protein expression are in italic and boldfaced characters;

figure 9 shows the Southern blot of total nucleic acids extracts from wild-type *N. benthamiana* protoplasts cotransfected with a TYLCSV infectious clone (pTOM6), along with the plasmid expressing the mutated Rep protein indicated above each column;

figure 10 shows the quantitative analysis of the TYLCSV replication in wild-type *N. benthamiana* protoplasts cotransfected with pTOM6 plasmid along with the plasmid expressing the mutated Rep protein;

figure 11 shows the scheme of pTOM130 plasmid used for obtaining Rep-130-expressing transgenic plants. LB and RB mean left- and right-border respectively; pE35S represents the duplicated *Cauliflower Mosaic Virus* 35S promoter; Rep-130 (SEQ ID No 8) is the sequence encoding the Rep-130 protein (SEQ ID No 9); t35S is the *Cauliflower*

Mosaic Virus 35S terminator; tNOS is the terminator of the gene encoding the nopaline synthase; nptII is the sequence encoding the neomycin phosphotransferase; pNOS is the promoter of the gene for the nopaline synthase; Kan is the gene for the kanamycin resistance;

5 figure 12 shows the analysis of the expression of Rep-130 protein (SEQ ID No 9) in transgenic *N. benthamiana* plants transformed with pTOM130 (lines 300-309);

 figure 13 shows the analysis of the TYLCSV replication in wild-type (wt) and transgenic *N. benthamiana* protoplasts expressing either the
10 Rep-130 protein (SEQ ID No 9) (lines 300, 301, 303) or the Rep-210 protein (102.22).

 figure 14 shows the analysis of the expression of Rep-130 protein (SEQ ID No 9) in transgenic *L. esculentum* plants transformed with pTOM130 (lines 402, 403, 406, 411, 413, 416, 417). A protein extract from
15 a transgenic Rep-130 expressing *N. benthamiana* (line 303) was used as positive control;

 figure 15 shows the comparison between a *L. esculentum* transgenic plant transformed with pTOM130 (line 406) expressing Rep-130 and a non-transformed wild-type plant;

20 figure 16 A and B shows two examples of synthetic sequences encoding Rep-210 (SEQ ID No 2, SEQ ID No 4). The alignment between the wild-type nucleotide sequence encoding TYLCSV Rep-210 protein (Seq_cod_Rep210_wild_type, on the top; SEQ ID No 1) and the synthetic nucleotide sequence modified so as to be an ineffective target of the virus-
25 induced post-transcriptional silencing is shown (Seq_cod_Rep210_silencing_minus, in the bottom; SEQ ID No 2, SEQ ID No 4). In the synthetic sequences, the mutated nucleotides with respect to the wild-type sequence are shaded;

 figure 17 shows the analysis of transient expression, by
30 agroinfiltration into *N. benthamiana* leaves, of Rep-210 protein encoded by the plasmid wild-type gene pTOM102(C4 -) and by the synthetic gene Rep-210 silencing minus B (SEQ ID No 4), (plasmid pTOM102 Syn);

 figure 18 shows a transient assay for the inhibition of viral replication through co-agroinfiltration of an *A. tumefaciens* strain
35 containing the TYLCSV infectious clone along with *A. tumefaciens* strains containing: a) the pTOM102(C4 -) plasmid expressing the wild-type gene for Rep-210 SEQ ID No 1 (Brunetti et al., 2001), lines 1-3; b) the

pTOM102Syn plasmid expressing the synthetic gene, for Rep-210 (Rep-210 silencing minus B; SEQ ID No 4) lines 4-6; c) the empty cloning plasmid pBIN19 lines 7-9;

5 figure 19 shows the analysis of the expression of Rep-210 protein in transgenic *N. benthamiana* plants transformed with pTOM102Syn plasmid containing the synthetic gene for Rep-210, Rep-210-silencing minus B, SEQ ID No 4 (lines 506, 508A and 508B). A protein extract from Rep-210-expressing transgenic tomato plant was used as positive control;

10 figure 20 shows the analysis of the expression of Rep-210 protein in transgenic *N. benthamiana* plants transformed with pTOM 102 (line 102.22, Noris et al., 1996) or with pTOM 102Syn (line 506) after TYLCSV agroinoculation. Analysis has been performed before (0wpi) and five weeks after (5wpi) TYLCSV agroinoculation;

15 figure 21 shows the analysis of the infection by "dot-blot" assay at 2, 3, 4, 5 wpi (where wpi means the number of weeks after the agroinoculation) on *N. benthamiana* wild-type (WT) or transgenic plants transformed with either pTOM 102 (line 102.22, Noris et al., 1996) or pTOM 102Syn (line 506);

20 figure 22 shows an example of synthetic sequence encoding CP. The alignment between the wild-type nucleotide sequence encoding TYLCSV CP (TYLCSV CP, on the top; SEQ ID No 12) and the synthetic nucleotide sequence modified so as to be an extremely ineffective or non target of the virus-induced post-transcriptional degradation (TYLCSV CP silencing minus, in the bottom; SEQ ID No 6) is shown. In the synthetic
25 sequence, the mutated nucleotides with respect to the wild-type sequence are shaded.

EXAMPLE 1: Identification of regions of the TYLCSV genome under-represented in the siRNAs population.

30 In a natural infection by TYLCSV of wild-type plants, the viral sequences transcribed by both strands of TYLCSV genome are target of post-transcriptional gene silencing as pointed out by the presence of siRNAs homologous to different portions of the genome (figure 5). In figure 5 is shown the Northern blot of total RNAs extracted from the tomato wild-type plants infected by TYLCSV (samples 1-4) and non-infected control (sample C). Probe and the restriction sites used are indicated aside each
35 panel. Also the estimated sizes of siRNAs are set forth.

In order to evaluate if some regions of the TYLCSV genome constitute a target of post-transcriptional gene silencing less effective than others, it was performed a systematic study of the siRNA distribution with respect to their position on the viral genome. Therefore the TYLCSV genome has been divided in nine contiguous fragments, each of about three hundred base pairs (as drawn in figure 6), obtained by PCR with specific oligonucleotides. The same amount of such fragments has been transferred on nylon filter after agarose gel electrophoresis. Quantification of PCR fragments loaded on agarose gel has been performed by software Aida. The siRNAs produced by a TYLCSV-infected tomato plant have been purified starting from the total RNAs, terminally labelled and used as probe (Szitty et al., 2002) on the filter containing several regions of TYLCSV genome. The different intensity of the hybridization signals, referred to a same amount of loaded fragment, has been assessed through the TYPHOON apparatus (Amersham-Pharmacia). So a different distribution of the siRNAs with respect to the several regions of the viral genome has been detected (fig.6).

EXAMPLE 2: Identification of a region of TYLCSV C1 gene under-represented in the si RNAs population.

In order to identify a region of the TYLCSV C1 gene under-represented in the siRNAs population, total RNAs (Brunetti et al., 1997) both from healthy and TYLCSV-infected tomato plants have been extracted.

Thirty micrograms of such RNAs have been submitted to 8% denaturing polyacrylamide gel electrophoresis and transferred by capillarity on nylon filter through Northern blot. Two identical replicas have been produced and for each it has been carried out an hybridization with probes corresponding to different portions of the 5' region of the C1 gene, as shown in figure 7. One filter has been hybridized with a probe derived from the 5' portion of C1 gene comprising 42 nt of non-translated leader sequence and the first 630 nucleotides of C1 gene (about 3/5 of C1 gene) (probe A) and the other filter with a probe derived from the 5' portion of C1 gene comprising 42 nt of non-translated leader sequence and the first 390 nucleotides of C1 gene (probe B).

In order to quantitatively compare the results obtained by the two different probes (deriving from two independent labelling), scalar amounts of a same 40mer oligonucleotide complementary to both probes

have been loaded on both replicas. Columns 100 and 50 correspond to 100 and 50 picograms of such oligonucleotide, respectively. The panels showing the oligonucleotide migration have been set close to the respective panels containing the siRNAs but their position in the figure
5 doesn't correspond to the position on gel, because the oligonucleotide and the siRNAs have different molecular weights.

Both probes after *in vitro* transcription have been submitted to alkaline hydrolysis (Cox et al., 1984) in order to obtain from them fragments with an average length of 75 nucleotides.

10 The hybridizations have been performed for 16 hours at 39°C in the buffer described by Dalmay et al., 2000. After hybridization the filters have been washed in 2X SSC, 0,2% SDS twice for 10 minutes at 40°C, twice for 10 minutes at 45°C and once for 10 minutes at 50°C.

It is remarkable how the proximal 5' region of the C1 gene in the
15 siRNAs population is under-represented. Particularly, the quantitative analysis of the results performed through the TYPHOON apparatus (Amersham-Pharmacia) revealed that the siRNAs corresponding to this 5' region are about 25% (probe B) with respect to those corresponding to the region extended up to nucleotides encoding the 210 aminoacid (probe A).
20 Said 5' region constitutes therefore an ineffective target for the virus-induced post-transcriptional gene silencing.

These results have been confirmed using the method described in example 1, i.e., where PCR fragments corresponding to the two different regions of the C1 gene were hybridised with the population of
25 siRNA extracted from tomato plants infected by TYLCSV.

EXAMPLE 3: *Construction of a polynucleotide sequence of the C1 gene 5' portion encoding the truncated Rep.*

As previously pointed out (Brunetti et al., 1997), the Rep-210 transgenic plants show a not long lasting resistance and an altered
30 phenotype.

As can be noticed in figure 1, the C4 gene is nested in the truncated C1 gene in a different reading frame.

It is shown that the transgenic expression of geminivirus C4 gene induces phenotype alterations (Krake et al., 1998).

35 Therefore, it has been designed several truncated C1 constructs unable to express C4 ORF.

In order to obtain C4 (-) mutants, a stop codon has been introduced in the C4 sequence through the introduction of two point mutations. Particularly, referring to the pTOM130 sequence set forth in figure 8 (SEQ ID No 8), the mutation at nucleotide 233 consists of a
5 trasversion from C to G that converts the TCA codon (encoding serine) of the reading frame encoding C4 in TGA (opal). In addition, the mutation at nucleotide 231 consists of a transition from C to T that restores in the reading frame encoding C1 a leucine codon (CTC becomes TTG, more represented in plant).

10 Thereby the translation of the C4 protein is interrupted after only 10 aminoacids, while the aminoacid sequence of the C1 protein remains unchanged. The two introduced mutations have been chosen among many possible mutations based on the criterion to generate a "strong" stop codon in the C4 reading frame, maintaining in the C1 reading
15 frame a leucine codon compatible with codon usage in plants.

Mutagenesis has been performed by PCR with the following mutated oligonucleotides:

C4 plus.primers (SEQ ID No 10): 5'-CT CAT CTC CAT ATT TTG
ATC CAA TTC GAA G-3'

20 C4 minus.primers (SEQ ID No 11): 5'-C TTC GAA TTG GAT CAA AAT ATG GAG ATG AG-3' (2419-2448 in TYLCV - Kheyr-Pour et al., 1991)

Each of the two mutated primers has been used along with an external primer in two separate PCR reactions using pGEM102 as
25 template (Brunetti et al., 2001).

Particularly, the external oligonucleotides are Rev and Univ (M13/pUC sequencing primer n.1233 and 1224). From the reaction performed with Univ/C4plus it has been obtained a 537 bp fragment, while from the reaction with Rev/C4minus a 351 bp fragment.

30 The obtained products have been used as templates for a following amplification reaction carried out using two external primers.

The obtained PCR product has been digested with EcoRI and BamHI restriction enzymes and cloned into the corresponding sites of pJIT60, thus obtaining pJITR210. In both cases it has been carried out the
35 sequencing to verify clones.

EXAMPLE 4: *Identification of the minimal 5' region of TYLCSV C1 gene that when expressed in plant cells is able to inhibit viral replication.*

In order to define the minimal 5' terminal region of C1 gene able to confer resistance against TYLCSV, a series of 3'-terminal deletion mutants of C1 gene was cloned in pJIT60 expression vector, resulting in a pJTR series.

The viral sequences have been amplified by PCR with Pfu DNA polymerase (Stratagene), using specific primers containing restriction sites at the ends.

The previously described pJTR210 plasmid, which encodes Rep-210, and contains a stop codon for the internal C4 protein, has been used as template. The fragments obtained by amplification reactions have been digested with BamHI and EcoRI enzymes and cloned in the corresponding sites of pJIT60 resulting in the pJTR series.

All final clones have been sequenced in order to confirm the amplification fidelity and vector-insert junctions. The length and the precise positions of every amplified sequence are set forth in table 6.

The ability of each Rep deletion mutant to confer resistance against TYLCSV has been evaluated through cotransfection assays of *N. benthamiana* wild-type protoplasts with a TYLCSV infectious clone (pTOM6) along with each mutant, and following analysed for the replication level of the viral genome through Southern blot. The obtained results are set forth in figures 9 and 10.

Table 6

pJTR210a	42 bp UTR + truncated C1 ORF (630 nt) containing C4 encoding region	1985-2656 (1)
pJTR210	42 bp UTR + truncated C1 ORF (630 nt)	1985-2656 (1)
pJTR181	42 bp UTR + truncated C1 ORF (543 nt)	2072-2656 (1)
pJTR156	42 bp UTR + truncated C1 ORF (468 nt)	2147-2656 (1)
pJTR130	42 bp UTR + truncated C1 ORF (390 nt)	2225-2656 (1)
pJTR120	42 bp UTR + truncated C1 ORF (360 nt)	2255-2656 (1)

25

pJTR80	42 bp UTR + truncated C1 ORF (240 nt)	2375-2656 (1)
pJTR54	42 bp UTR + truncated C1 ORF (162 nt)	2453-2656 (1)

(1) nucleotide numbering of the TYLCSV genome are according to Kheyr-Pour et al. 1991.

The protoplast cotransfection, total nucleic acid extraction and Southern analysis have been performed according to already described methods (Brunetti et al. 2001).

Total nucleic acids extracts from each protoplast sample have been analysed through Southern blot with a digoxigenin-labelled RNA probe corresponding to the sequence encoding Rep-210, and the pGEM-P plasmid used as control. In particular figure 9 represents a Southern blot of total nucleic acids, where scDNA and ssDNA mean supercoiled and single strand DNA of TYLCSV, respectively.

For an accurate quantitative analysis of the effect of the expression of several truncated forms of Rep on the replication of TYLCSV genome, a Southern analysis has been performed with a ³²P-labelled DNA probe corresponding to the region encoding the first 54 N-terminal aminoacids of Rep and the radioactivity level corresponding to each band detected on filter has been evaluated, through analysis with the Istant Imager apparatus (Canberra, Packard).

Each mutated construct has been assayed in duplicate, in three independent experiments and the value set forth in figure 10 represents the average of two or three cotransfections of the three independent experiments.

The level of TYLCSV replication in the cotrasfection experiments performed with pTOM6 along with pGEM-P control plasmid was considered equal to 100%.

Particularly, figure 10 shows a quantitative analysis, the white and black bars of the histograms represents the amounts of supercoiled and single strand DNA, respectively; error bars indicates the mean standard deviation.

As pointed out by observing figures 9 and 10, the first 130 N-terminal aminoacids of the Rep protein are enough to inhibit almost completely viral replication, while the expression of the first 120 N-terminal aminoacids has no influence.

EXAMPLE 5: *Production of N. benthamiana transgenic plants expressing Rep-130.*

The analysis of the ability to inhibit TYLCSV replication by the Rep mutants assessed through transient expression in protoplasts, has revealed that the shortest mutant still effective encodes Rep-130 (SEQ ID No 9) as described in the preceding example.

Also it has been previously revealed in other examples that the proximal 5' portion of C1 gene encoding Rep-130 is a less effective target of post-transcriptional gene silencing compared to sequence encoding Rep-210.

Therefore it has been obtained *N. benthamiana* transgenic plants expressing Rep-130. For this purpose, the pTOM130 plasmid represented in figure 11 has been obtained, by cloning KpnI-BglII fragment of pJTR130 into the KpnI-BamHI sites of pBIN19.

N. benthamiana has been transformed with the *A. tumefaciens* pGV2260 C58 strain containing pTOM130 plasmid and plants resistant to kanamycin have been regenerated as described (Noris et al., 1996).

The primary transformants have been analysed for the presence of transgene by PCR analysis and for the expression of Rep-130 protein through Western blot, as shown in figure 12.

The protein extracts obtained from transgenic (300-309) or wild-type control (wt) plants have been analysed by Western blot using an anti-TYLCSV Rep rabbit polyclonal primary antibody as described (Noris et al., 1996).

EXAMPLE 6: *Plant cells stably transformed with the pTOM130 construct and expressing Rep-130 inhibit TYLCSV replication.*

In order to early evaluate the resistance conferred by Rep-130, protoplasts isolated from several primary transgenic Rep-130 expressing *N. benthamiana* plants were transfected with a TYLCSV infectious clone (pTOM6).

Transgenic lines have been chosen for their high Rep-130 expression, as revealed by Western blot analysis (figure 12).

The level of TYLCSV replication in such transgenic protoplasts has been compared with that observed in *N. benthamiana* wild-type and in transgenic protoplasts expressing Rep-210 (line 102.22).

Particularly, figure 13 shows the analysis of TYLCSV replication in wild-type (wt) and transgenic *N. benthamiana* protoplasts expressing

either Rep-130 (SEQ ID No 9) (lines 300, 301, 303), or Rep-210 (line 102.22). Total nucleic acids extracts from several protoplast samples have been analysed by Southern blot using a digoxigenin-labelled RNA probe corresponding to the Rep-210 transcript.

5 In order to compare the level of TYLCSV replication in the Rep-130 transgenic protoplasts with that observed in wild-type protoplasts, the total nucleic acids extracted from wild-type protoplasts have been also loaded following 1:10 and 1:50 dilutions, as shown in figure 13.

10 EXAMPLE 7: Production of tomato transgenic plants expressing Rep-130

The analysis of the ability to inhibit TYLCSV replication by Rep mutants, assessed by transient expression in protoplasts, has revealed that the shortest mutant still effective encodes Rep-130, as described in the previous example.

15 As previously shown, the proximal 5' portion of C1 gene encoding Rep-130 is an ineffective target of post-transcriptional gene silencing compared with the sequence encoding Rep-210.

20 Therefore it has been carried out the production of transgenic tomato plants (*Lycopersicon esculentum* cv. Moneymaker) expressing Rep-130.

The tomato has been transformed using *A. tumefaciens* pGV2260 C58 strain containing pTOM130 plasmid (figure 11) and kanamycin-resistant plants were regenerated as described (Brunetti et al. 1997).

25 The primary transformants have been analysed for the presence of the transgene by PCR analysis and for the expression of Rep-130 protein by Western blot (figure 14).

30 The protein extracts obtained from transgenic (lines 400) or wild-type control (wt) plants have been analysed by Western blot using an anti-TYLCSV Rep polyclonal rabbit primary antibody as described (Noris et al. 1996).

All transgenic tomato plants expressing Rep-130 protein are phenotypically impossible to distinguish from wild-type plants (figure 15).

35 EXAMPLE 8: Demonstration of long-lasting resistance against TYLCSV in plants transgenic for the pTOM130 construct expressing Rep-130

In order to assess the lasting of the resistance against TYLCSV conferred by Rep-130 expression, the *N. benthamiana* R1 transgenic plants expressing Rep-130 have been agroinoculated with the *A. tumefaciens* LBA4404 strain containing the TYLCSV infectious clone.

5 As previously reported, the viral delivery through agroinoculation, used to assay the resistance and evaluate stability over time, corresponds to high or very high viral pressure conditions.

Infection of plants has been assessed at weekly intervals by a "tissue printing" assay, using a digoxigenin-labelled probe specific for the coat protein gene.

10 The results in table 7 show that, unlike the results described in table 2 concerning transgenic plants expressing Rep-210, transgenic *N. benthamiana* plants expressing Rep-130 protein show a long-lasting resistance when agroinoculated with TYLCSV. This can be deduced by comparison of the resistant plants at 2 and 6 weeks following inoculation.

Table 7

Plants	Time (weeks)	N° infected plants/inoculated plants	% infected plants / inoculated plants
Rep-130	2	0/10	0
	4	0/10	0
	6	0/10	0
Wild-type	2	9/10	90
	4	10/10	100

In addition, it was assessed the stability of the resistance against TYLCSV conferred by Rep-130 expression in transgenic R2 tomato plants.

20 The plants have been agroinoculated with the *A. tumefaciens* C58C1+pCH32 strain containing TYLCSV infectious clone. The agroinoculation conditions used for assaying the resistance and evaluating stability thereof over time, correspond to high or very high viral pressure conditions.

The infection has been assessed at intervals of one or two weeks through dot-blot assay, using a radioactively labelled probe specific for the coat protein gene.

- 5 The results in table 8 point out that tomato transgenic plants expressing Rep-130 show a long lasting resistance when agroinoculated with an TYLCSV infectious clone. This can be deduced from the comparison of the resistant plants at 3 and 12 weeks after inoculation.

Table 8

Plants	Time (weeks)	N° infected plants/inoculated plants	% infected plants / inoculated plants
Rep-130	3	0/9	0
	4	0/9	0
	5	0/9	0
	6	0/9	0
	7	0/9	0
	8	0/9	0
	10	0/9	0
	12	0/9	0
segregating Rep-130	3	3/3	100
Wild-type	3	4/4	100

- 10 Therefore the resistance is associated to the presence of Rep-130 protein (SEQ ID No 9) and to the ability of TYLCSV-inoculated transgenic plant to stably express Rep-130, because the sequence encoding it is an ineffective target of virus-induced post-transcriptional gene silencing and Rep, even if further mutated, maintains its ability to confer virus resistance.

- 15 **EXAMPLE 9:** *Construction of a synthetic polynucleotide sequence, modified in order not to be or to be an extremely ineffective target of the post-transcriptional gene silencing induced by the infecting virus, encoding TYLCSV Rep-210 protein.*

- 20 In order to achieve the long lasting expression of TYLCSV Rep-210 protein in transgenic plants, it has been produced, employing the method according to the present invention, a synthetic polynucleotide sequence, able to encode for Rep-210 protein, that is not or is an ineffective target of the post-transcriptional gene silencing induced by the infecting virus.

In addition, the following criteria have been followed:

- the synthetic polynucleotide sequence is not able to encode the C4 protein, having in positions 231 and 233 the same mutations shown in figure 8, as described for Rep-130 (SEQ ID No 8; SEQ ID No 9).
- 5 - the introduced mutations are all silent, namely the protein product encoded by the synthetic polynucleotide sequence matches that encoded by the viral wild-type sequence;
- the mutations were introduced according to the frequency of codon usage in the tomato genes; particularly whenever possible the more frequently used codon in tomato was selected;
- 10 - the introduced mutations have been all checked to exclude the possible formation of sequences having a particular function, such as for example polyadenylation or splicing signals, also cryptic.

Following the above described criteria, two synthetic sequences encoding Rep-210 have been designed (figure 16 A and B, SEQ ID No 2, SEQ ID No 3, SEQ ID No 4, SEQ ID No 5).

A non-translated leader sequence at the 5' and a stop codon at the 3' have been added to the sequence of the synthetic Rep-210 silencing minus B gene (SEQ ID No 4).

20 Particularly, the polynucleotide sequence containing in the 5'-3' order the non-translated leader sequence, the synthetic sequence encoding Rep-210 (figure 16 B; SEQ ID No 4, SEQ ID No 5) and the stop codon has been assembled by PCR starting from oligonucleotides (Prodromou and Laurence, 1992; Stemmer et al., 1995), using a

25 thermostable DNA polymerase with "proof reading" correction activity (Pfu DNA Polymerase, Stratagene and/or Pfx DNA Polymerase, Invitrogen).

The synthetic gene has been subsequently cloned in pJT60 plasmid under the transcriptional control of 35S promoter of the *Cauliflower mosaic virus* (CaMV) and the transcription termination sequences of the CaMV 35S, producing the pJT60Syn. Then the cassette containing in the 5'-3' order: 35S promoter, Rep-210 synthetic gene, 35S terminator, has been removed from pJT60Syn plasmid by restriction with *KpnI-BglII* and cloned in the *KpnI-BamHI* sites of the binary plasmid pBIN19 generating pTOM102Syn.

35 EXAMPLE 10: *Evaluation of the inhibition of viral replication by Rep-210 synthetic gene.*

The correct expression of Rep-210 protein, encoded by the synthetic gene, has been checked through agroinfiltration of *N. benthamiana* leaves, with *A. tumefaciens* C58C1/pCH32 transformed with pTOM102Syn. The strain C58C1/pCH32 transformed with pTOM102 (C4 -) has been used as a positive control, while as negative control the strain C58C1/pCH32 transformed with the binary plasmid pBIN19 was used. Western blot analysis (figure 17) shows the expression of Rep-210 protein encoded by the synthetic gene.

In order to assess the ability of the Rep-210 protein, encoded by pTOM102Syn, to inhibit TYLCSV replication, a transient co-agroinfiltration assay has been carried out. *N. benthamiana* leaves have been co-agroinfiltrated with *A. tumefaciens* C58C1/pCH32 strain containing the TYLCSV infectious clone (pTOM6) along with the *A. tumefaciens* C58C1/pCH32 strain containing: a) pTOM102Syn plasmid; b) pTOM102 (C4-) plasmid; c) pBIN19 binary plasmid. The TYLCSV replication has been assessed through Southern analysis of the total nucleic acids extracted from the co-agroinfiltrated tissues 72 hours after the infiltration. This analysis has pointed out that Rep-210 protein expressed by the synthetic gene (pTOM102Syn) and by pTOM102(C4 -) wild-type gene, inhibits TYLCSV replication in a similar manner (figure 18).

EXAMPLE 11: *Production of transgenic N. benthamiana plants expressing the synthetic gene for Rep-210.*

In order to obtain transgenic *N. benthamiana* plants expressing the synthetic gene for the Rep-210, *N. benthamiana* leaf-discs have been transformed using the *A. tumefaciens* LBA 4404 strain containing pTOM102Syn plasmid and the kanamycin-resistant plants have been regenerated as described (Noris et al. 1996).

The primary transformants have been analyzed for the expression of Rep-210 protein by Western blot analysis. Four primary transformants, 506, 508, 517 and 537 lines accumulating intermediate levels of Rep-210 have been selected for further studies. Figure 19 shows western analysis of proteins extracted from the 506, 508A and 508B plants.

EXAMPLE 12: *Stability of the resistance in N. benthamiana transgenic plants for Rep-210 synthetic gene.*

The authors have previously shown (Noris et al. 1996; Brunetti et al. 1997) that there is a direct correlation between the amounts of Rep-

210 protein produced by the transgenic plants and resistance against TYLCSV. Transgenic plants transformed with the pTOM102 construct accumulating intermediate levels of Rep-210 protein are susceptible to viral infection, like non-transformed plants (Noris et al. 1996 and unpublished data). The low level of Rep-210 protein in these plants is not enough to completely inhibit viral replication, thus allowing the establishment of an early virus-induced post-transcriptional silencing leading to a drastic reduction in Rep-210 protein accumulation which causes lack of resistance.

In order to assess if Rep-210 protein encoded by the synthetic gene is not or is an ineffective target of virus-induced post-transcriptional gene silencing and therefore to control over time the viral infection, line 102.22 transgenic plants (R3) and line 506 transgenic plants (R0) expressing similar amount of Rep-210 (figure 20, 0 wpi), have been agro-inoculated with the TYLCSV and analysed by dot blot at week intervals for the accumulation of the TYLCSV. As expected (Noris et al., 1996) the transgenic R3 line 102.22 plants (figure 21, 5-8) that accumulate intermediate levels of Rep-210 protein are susceptible as the non-transformed plants (figure 21, 1-4). As shown by dot blot analysis (figure 21, 9-12) the transgenic plants for the synthetic construct (R0 line 506) are resistant to viral infection, accumulating only limited amounts of virus. Interestingly and according to virus inability to post-transcriptionally silence effectively the synthetic gene, Rep-210 was still accumulating 5 weeks after inoculum (figure 20, 5 wpi) and inhibiting over time TYLCSV replication (figure 21, 9-12).

The results described in the examples point out that it is possible to obtain a long lasting resistance against geminiviruses by expressing in plant a transgene consisting of a pathogen-derived polynucleotide sequence, if the latter is suitably selected or modified in order not to be a target or to be an ineffective target of the post-transcriptional gene silencing by the infecting virus.

EXAMPLE 13: *Construction of a synthetic polynucleotide sequence modified in order not to be a target or to be an extremely ineffective target, of the post-transcriptional degradation induced by the infecting virus, encoding the TYLCSV capsid protein.*

As above reported, the transgenic expression of the TYLCV capsid protein in a interspecific tomato hybrid (*Lycopersicon esculentum* X

L. pennellii) confers a partial resistance against viral infection (Kunik et al., 1994). Also in this case, the resistance mediated by the expression of the capsid protein is not long lasting.

5 In order to obtain a stable expression of the TYLCSV capsid protein (CP) by transgenic plants, it has been produced a synthetic polynucleotide sequence, able to encode the CP, which results in an ineffective target of virus-induced post-transcriptional gene silencing.

10 The synthetic polynucleotide sequence has been designed so as to satisfy the requisite not to be or to be an extremely ineffective target of virus-induced post-transcriptional gene silencing employing the method according to the present invention.

In addition, the following criterions have been followed:

- the introduced mutations are all silent, namely the protein product encoded by the synthetic polynucleotide sequence is exactly matching that encoded by the viral wild-type sequence;
- the introduced mutations consider the frequency of codon usage in the tomato genes; particularly whenever possible the codon more frequently used in tomato is selected;
- the introduced mutations have been all checked to exclude the possible formation of sequences having a particular function, such as for example polyadenylation signals or splicing signals, also cryptic.

20 Following the above described criterions a synthetic sequence encoding CP has been designed (SEQ ID No 12) (figure 22, TYLCSV CP silencing minus, SEQ ID No 6, SEQ ID No7).

25

BIBLIOGRAPHY

- Abel PP, Nelson RS, De B, Hoffmann N, Rogers SG, Fraley RT; Beachy RN. 1986. Delay of disease development in transgenic plants that express the tobacco mosaic virus coat protein gene. Science 232:738-43.
- Baulcombe DC 1996. Mechanisms of Pathogen-Derived Resistance to Viruses in Transgenic Plants. 8:1833-1844.
- Beachy, R. 1997. Mechanisms and applications of pathogen-derived resistance in transgenic plants. Current Opinion in Biotechnology 8:215-220.

35

- Briddon, R.W. and P.G. Markham. 1995. *Geminiviridae*, Sixth report of international committee on taxonomy of viruses. Springer-Verlag, New York, N.Y.
- Brunetti, A., R. Tavazza, E. Noris, A. Lucoli, G. P. Accotto, and M. Tavazza, 2001, J Virol. 75:10573-81.
- Cahill, M., K. Gorman, S. Kay and I. Denliolm.1996. Baseline determination and detection of resistance to imidacloprid in Bemisia tabaci (Homoptera: Aleyrodidae) Bulletin of Entomology.Research 86:343-349.
- Cox, K. H., D. V. DeLeon, L. M. Angerer, and R. C. Angerer. 1984. Detection of mrnas in sea urchin embryos by in situ hybridization using asymmetric RNA probes. Dev Biol 101:485-502
- Czosnek, H., and H. Laterrot. 1997. A world survey of tomato yellow leaf curl viruses. Archives of Virology 142:1391-1406.
- Dalmay, T., A. Hamilton, E. Mueller, and D. C. Baulcombe. 2000. Potato virus X amplicons in arabidopsis mediate genetic and epigenetic gene silencing. Plant Cell 12:369-379.
- English J., Mueller E., and Baulcombe D.C. 1996. Suppression of Virus Accumulation in Transgenic Plants Exhibiting Silencing of Nuclear Genes. Plant Cell. 8: 179-188.
- Fauquet CM, Bisaro DM, Briddon RW, Brown JK, Harrison BD, Rybicki EP, Stenger DC, Stanley J. 2003. Revision of taxonomic criteria for species demarcation in the family *Geminiviridae*, and an updated list of *begomovirus* species. Arch Virol. 148:405-21.
- Hanley-Bowdoin, L., S.B. Settagle, B.M. Orozco, S. Nagar and D. Robertson. 1999. *Geminivirus*: models for plant DNA replication, transcription, and cell cycle regulation. Critical Reviews in Plant Sciences, 18:71-106.
- Kheyr-Pour A., M. Bendahmane, V. Matzeit, G.P. Accotto, S. Crespi and B. Gronenborn. 1991. Tomato yellow leaf curl virus from Sardinia is a whitefly-transmitted monopartite *Geminivirus*. Nucleic Acids Research 19:6763-6769. Picò, B., M.J. Diez and F. Nuez. 1996. Viral disease causing economic losses to the tomato crop. II. The tomato yellow leaf curl virus review. Scientia Horticulturæ 67:151-196.
- Krake, L.R., M.A Rezaian, and I.B. Dry. 1998. Expression of the tomato leaf curl *Geminivirus* C4 gene produces viruslike

symptoms in transgenic plants *Molecular Plant Microbe Interactions* 11:413-417.

- 5 - Kunik, T., R. Salomon, D. Zamir, N. Navot, M. Zeidan, I. Michelson, Y. Gafni and H. Czosnek 1994. Transgenic tomato plants expressing the tomato yellow leaf curl virus capsid protein are resistant to the virus. *Bio/technology* 12:500-504.
- 10 - Noris, E., G. P. Accotto, R. Tavazza, A. Brunetti, S. Crespi, and M. Tavazza. 1996. Resistance to tomato yellow leaf curl *Geminivirus* in *Nicotiana benthamiana* plants transformed with a truncated viral C1 gene [published erratum appears in *Virology* 1997 Jan 20;227(2):519]. *Virology* 224:130-8.
- 15 - Polston, J. E. and P. K. Anderson. 1997. The emergence of whitefly-transmitted gemini viruses in tomato in the Western hemisphere. *Plant Disease* 81:1358-1369.
- 20 - Prodromou, C. and L.H. Pearl. 1992. Recursive PCR: a novel technique for total gene synthesis. *Protein Engineering*, 5:827-829.
- 25 - Sangaré, A., D. Deng, C.M. Fauquet and R. Beachy. 1999. Resistance to African cassava mosaic virus conferred by a mutant of the putative NTP-binding domain of the Rep gene (AC1) in *Nicotiana benthamiana*. *Molecular Biology Reports* 5:95-102.
- 30 - Stemmer, W.P.C., A. Cramer, K.D. Ha, T.M. Brennan, H.L. Heyneker. 1995. Single-step assembly of a gene and entire plasmid from large numbers of oligodeoxynucleotides. *Gene*, 164:49-53.
- 35 - Szittya, G., D. Silhavy, A. Molnar, C. Hornyik, J. Burgyn. 2002. Short Defective Interfering RNAs of Tombusviruses Are Not Targeted but Trigger Post-Transcriptional Gene Silencing against Their Helper Virus. *The Plant cell*, Vol. 14, 359-372.
- Szittya, G., D. Silhavy, A. Molnar, Z. Havelda, A. Lovas, L. Lakatos, Z. Banfalvi and J. Burgyn. 2003. Low temperature inhibits RNA silencing-mediated defence by the control of siRNA generation. *EMBO Journal* 22:1-8.
- Tavazza, M., A. Lucoli. 1993. Approcci Molecolari per la Resistenza alle Virosi. In "Miglioramento Genetico per Resistenza a Patogeni e Parassiti." *Fondamenti Teorici e Pratici*. Buiatti M.,

Crinò P., Porta-Puglia F., Saccardo F., Sonnino A., Surico G.
(Editori). Ed. Agricole Bologna. 191-214.

Voinnet, O. 2001. RNA silencing as a plant immune system
against viruses. Trends in Genetics 17:449-459.

5 Waterhouse, P.M., M.B. Wang and T. Lough. 2001. Gene
silencing as an adaptive defence against viruses. Nature 411:834-
842.

Williams, L., T. J. Dennehy and J. C. Palumbo 1996. Development
of a resistance management program for imidacloprid"Proc.
10 Beltwide Cotton Conferences pp. 752-755.

Zaitlin M. and Palukaitis P. 2000. Advances in understanding plant
viruses and virus diseases. Annual review of phytopathology 38:
117-143.

15

CLAIMS

1. Polynucleotide sequence encoding a geminivirus-derived aminoacid sequence, said polynucleotide sequence being characterised in that is not or is an ineffective target of the viral post-transcriptional silencing and has:
- 5 a) an homology at nucleotidic level below or equal at 90% with respect to the corresponding gene sequence of the geminiviruses against which the resistance is required;
- 10 b) a continuous homology in the transcribed RNA, with respect to the corresponding gene sequence of the geminiviruses, below or equal to 17 nucleotides;
- 15 c) a maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of the geminiviruses not higher than 30 nucleotides
- said polynucleotide sequence being able to confer to the plants, tissues or plant cells therewith transformed, a lasting resistance against the geminiviruses.
2. Sequence according to claim 1, wherein the homology at the nucleotidic level with respect to the corresponding gene sequence of the geminivirus is below or equal to 80%.
- 20 3. Sequence according to claim 1, wherein the homology at the nucleotidic level with respect to the corresponding gene sequence of the geminivirus is below or equal to 70%.
- 25 4. Sequence according to each of the preceding claims, wherein the continuous homology in the transcribed RNA with respect to the gene sequence of the geminiviruses is below or equal to 8 nucleotides.
5. Sequence according to each of the preceding claims, wherein the continuous homology in the transcribed RNA with respect to the gene sequence of the geminiviruses is below or equal to 5 nucleotides.
- 30 6. Sequence according to each of the preceding claims, wherein the maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of the geminiviruses is not more than 20 nucleotides.
- 35 7. Sequence according to each of the preceding claims, wherein the maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of the geminiviruses is not more than 9 nucleotides.

8. Sequence according to each of the preceding claims, wherein the polynucleotide sequence has been mutated or it is a wild-type sequence selected from geminivirus so as to differ at the nucleotidic level with respect to the corresponding genomic sequence of the geminivirus against which a resistance is required according to each of the preceding claims.

9. Sequence according to each of the preceding claims, wherein the geminiviruses are selected from the group consisting of species of *Mastrevirus*, *Curtovirus*, *Begomovirus* and *Topocuvirus* and isolates thereof.

10. Sequence according to claim 9, wherein *Begomoviruses* species are selected from the group consisting of TYLCCNV, TYLCCGV, TYLCMaIV, TYLCSV, TYLCTHV, TYLCV, ACMV, BGMV, CaLCuV, ToCMoV, TGMV, ToGMoV, ToMHV, ToMoTV, ToMoV, ToRMV, ToSLCV, ToSRV, Cotton leaf curl (CLCrV, CLCuAV, CLCuGV, CLCuKV, CLCuMV, CLCuRV), East African cassava mosaic (EACMCV, EACMMV, EACMV, EACMZV), Potato yellow mosaic (PYMPV, PYMTV, PYMV), Squash leaf curl (SLCCNV, SLCV, SLCYV), Sweet potato leaf curl (SPLCGV, SPLCV), Tobacco leaf curl (TbLCJV, TbLCKoV, TbLCYNV, TbLCZV), Tomato leaf curl (ToLCBV, ToLCBDV, ToLCGV, ToLCKV, ToLCLV, ToLCMV, ToLCNDV, ToLCSLV, ToLCTWV, ToLCVV, ToLCV) and isolates thereof.

11. Sequence according to claim 9, wherein the species belonging to the genus *Mastrevirus*, *Curtovirus*, *Topocuvirus* are selected from the group consisting of WDV, MSV, SSV, BYDV, TYDV, BCTV and isolates thereof.

12. Sequence according to claim 1, wherein the gene sequence is selected from the group consisting of C1/AL1/AC1, C2/AL2/AC2, C3/AL3/AC3, C4/AL4/AC4, V1/AR1/AV1, V2/AR2/AV2, BC1/BL1 and BV1/BR1, belonging to the geminiviruses.

13. Sequence according to claim 12, wherein C1/AL1/AC1 gene sequence is from the geminiviruses, as defined according to claims 10 and 11.

14. Sequence according to claim 1, wherein the geminivirus aminoacidic sequence is a pathogen-derived protein able to confer resistance against the geminiviruses to the plants expressing the protein.

15. Sequence according to claim 14, wherein the protein is selected from the group consisting of capsid protein, replication associated

viral protein (Rep), proteins encoded by the genes C2/AL2/AC2, C3/AL3/AC3, C4/AL4/AC4, V2/AR2/AV2, BC1/BL1 and BV1/BR1.

16. Sequence according to claim 1, wherein the plants, tissues or cells thereof belong to the group consisting of tomato, pepper, tobacco,
5 squash, manioc, sweet potato, cotton, melon, potato, soybean, wine, corn, wheat, sugar cane, bean, beet.

17. Polynucleotide sequence encoding a geminivirus-derived aminoacid sequence, said polynucleotide sequence being characterised in that is not or is an ineffective target of the viral post-transcriptional silencing
10 and has homology even equal to 100% with respect to the sequence of the geminivirus against which a resistance is required and is suitably shortened to be under represented in the siRNAs population with respect to the original sequence.

18. Construct comprising an heterologous polynucleotide sequence containing in the 5'-3' direction:

a) a polynucleotide sequence acting as promoter in said plant or tissue or transformed cells;

b) a non translated polynucleotide sequence positioned 5' of the encoding region;

20 c) a polynucleotide sequence as defined according to claims from 1 to 17, a fragment or a variant thereof;

d) a sequence acting as transcription terminator, positioned 3' with respect to said polynucleotide sequence.

19. Expression vector comprising the construct as defined
25 according to claim 18.

20. Transgenic plant, tissue or plant cells thereof, comprising in their genome a polynucleotide sequence as defined according to claims from 1 to 17.

21. Progeny of the plants and plant tissues according to the
30 claim 20.

22. Seed comprising in its genome a polynucleotide sequence as defined according to claims from 1 to 17.

23. Method for the preparation of transgenic plants, plant tissue or cells thereof long lasting resistant against geminiviruses including the
35 following steps:

a) identification or selection of a viral gene sequence encoding an aminoacid sequence able to confer resistance against geminiviruses;

b) mutagenesis or choice of the viral gene sequence so as to make it an ineffective target of the post-transcriptional silencing induced by infecting geminivirus;

5 c) insertion of the geminivirus gene sequence mutated or chosen in the step b) in the plant, plant tissue or cell thereof using a construct as defined according to claim 18.

24. Method according to claim 23, wherein by mutagenesis the homology at the nucleotidic level with respect to the gene sequence of the geminivirus against which resistance is required is maintained below or
10 equal to 90 %, distributed in such way that the continuous homology in the transcribed RNA with respect to the corresponding gene sequence of the geminiviruses is below or equal to 17 nucleotides and the maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of the geminiviruses is not higher than 30
15 nucleotides.

25. Method according to anyone of the claims 23 and 24, wherein by mutagenesis an homology at the nucleotidic level with respect to the gene sequence of geminivirus is maintained below or equal to 80%.

26. Method according to anyone of the claims 23 and 24,
20 wherein by mutagenesis an homology at a nucleotidic level with respect to the gene sequence of geminivirus is maintained below or equal to 70%.

27. Method according to anyone of the claims from 23 to 26, wherein by mutagenesis a continuous homology at the nucleotidic level with respect to the gene sequence of geminivirus is maintained below or
25 equal to 8 nucleotides.

28. Method according to anyone of the claims from 23 to 27, wherein by mutagenesis a continuous homology at the nucleotidic level with respect to the gene sequence of geminivirus is maintained below or equal to 5 nucleotides .

30 29. Method according to anyone of the claims from 23 to 28, wherein the maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of geminiviruses is no more than 20 nucleotides.

30. Method according to anyone of the claims from 23 to 29,
35 wherein the maximum length of the sequence containing a single substitution with respect to the corresponding gene sequence of geminiviruses is no more than 9 nucleotides.

31. Method according to anyone of the claims from 23 to 30, wherein the mutagenesis consists of silent point mutations or deletions and/or insertions and/or substitutions.

5 32. Method according to anyone of the claims from 23 to 31, wherein the mutagenesis in step b) consists of deletions of the 5' or 3' regions of the viral gene sequence of step a) until the identification of the minimum region of said gene sequence that is under represented with respect to the sequence encoding the wild-type protein, in the population of the interfering siRNAs and that said truncated protein maintains the ability to confer resistance against geminiviruses.

10 33. Method according to claim 32, wherein the viral gene sequence of step a) is the C1/AL1/AC1 gene.

34. Method according to claim 32, wherein the C1/AL1/AC1 gene is a TYLCSV gene.

15 35. Method according to claim 32, wherein the aminoacid sequence is a truncated protein with respect to the viral wild-type protein.

36. Method according to anyone of the claims from 32 to 35 in which the viral gene sequence made not target or ineffective target of the post-transcriptional silencing is the SEQ ID No 8.

20 37. Method according to anyone of the claims from 32 to 36, wherein the truncated protein is Rep-130 (SEQ ID No 9).

38. Method according to anyone of the claims from 23 to 31, wherein the mutagenesis in step b) consists of silent point mutations of the viral gene sequence of step a) to maintain the ability of the aminoacid sequence, encoded by the same, to confer resistance against geminiviruses and not to be or to be an ineffective target of the post-transcriptional silencing.

25 39. Method according to claim 38, wherein the viral gene sequence of step a) is the V1/AR1/AV1 (CP) gene.

30 40. Method according to claim 39, wherein the V1/AR1/AV1 (CP) gene is a TYLCSV gene.

41. Method according to anyone of the claims from 38 to 40, wherein the viral gene sequence made not target or ineffective target of the post-transcriptional silencing is the SEQ ID No 8.

35 42. Method according to claim 38, wherein the viral gene sequence of step a) is C1/AL1/AC1 of TYLCSV.

42

43. Method according to claim 42, wherein the viral gene sequence made not target or ineffective target of the post-transcriptional silencing is the SEQ ID No 2 or the SEQ ID No 4.

ABSTRACT OF THE DISCLOSURE

The invention relates to a method for the production of transgenic plants durably resistant against Geminiviruses, wherein the transgene consists of a suitably modified polynucleotide sequence, derived by the pathogen.




Fig. 1

2/16




Fig. 2




Fig. 3

3/16




Fig. 4

4/16




Fig.5

5/16




Fig. 6

6/16




Fig. 7

7/16

SEQ ID No 8

```
GGATCCCCctggatactttgagtgtcccccgattcagaac 40
gacagcaaaaatgccaagatcaggtcgtttttagtatcaag 80
gctaaaaaattattttccttacatatcccaaattgtgatttaa 120
caaaagaaaatgcactttcccaaataacaaacctacaaac 160
accacaaacaaattattcatcaaaatttgcagagagaacta 200
catgaaaatggggaacctcatctccatatttggatccaat 240
tcgaaggaaaatacaattgtaccaatcaacgattcttcga 280
cctggtatccccaaccaggtcagcacatttccatccgaac 320
attcagggagctaaatcgagctccgacgtcaagtcctata 360
tcgacaaggacggagatgttcttgaatgggggtactttcca 400
gatcgacggacgatctgctaggggaggacaacagacagcc 440
tGAATTC 447
```

Fig. 8

8/16




Fig. 9




Fig. 10

9/16




Fig. 11

10/16




Fig. 12




Fig. 13

11/16




Fig. 14




Fig. 15

[illegible]

Fig. 16A

13/16

	10	20	30	40	50	60	
1	A	T	G	C	C	A	Rep-210 wild type (SEQ ID No 1)
1	A	T	G	C	C	A	Rep-210 silencing minus B (SEQ ID No 4)
	70	80	90	100	110	120	
61	T	G	T	G	A	T	Rep-210 wild type (SEQ ID No 1)
61	T	G	T	G	A	T	Rep-210 silencing minus B (SEQ ID No 4)
	130	140	150	160	170	180	
121	A	A	A	T	T	A	Rep-210 wild type (SEQ ID No 1)
121	A	A	A	T	T	A	Rep-210 silencing minus B (SEQ ID No 4)
	190	200	210	220	230	240	
181	C	T	C	A	T	C	Rep-210 wild type (SEQ ID No 1)
181	C	T	C	A	T	C	Rep-210 silencing minus B (SEQ ID No 4)
	250	260	270	280	290	300	
241	C	C	A	A	C	C	Rep-210 wild type (SEQ ID No 1)
241	C	C	A	A	C	C	Rep-210 silencing minus B (SEQ ID No 4)
	310	320	330	340	350	360	
301	A	A	G	T	C	A	Rep-210 wild type (SEQ ID No 1)
301	A	A	G	T	C	A	Rep-210 silencing minus B (SEQ ID No 4)
	370	380	390	400	410	420	
361	C	G	A	T	C	G	Rep-210 wild type (SEQ ID No 1)
361	C	G	A	T	C	G	Rep-210 silencing minus B (SEQ ID No 4)
	430	440	450	460	470	480	
421	G	G	A	G	T	C	Rep-210 wild type (SEQ ID No 1)
421	G	G	A	G	T	C	Rep-210 silencing minus B (SEQ ID No 4)
	490	500	510	520	530	540	
481	C	A	T	T	C	A	Rep-210 wild type (SEQ ID No 1)
481	C	A	T	T	C	A	Rep-210 silencing minus B (SEQ ID No 4)
	550	560	570	580	590	600	
541	G	T	T	C	T	C	Rep-210 wild type (SEQ ID No 1)
541	G	T	T	C	T	C	Rep-210 silencing minus B (SEQ ID No 4)
	610	620	630				
601	T	C	C	G	A	G	Rep-210 wild type (SEQ ID No 1)
601	T	C	C	G	A	G	Rep-210 silencing minus B (SEQ ID No 4)

Fig. 16B

14/16




Fig. 17




Fig. 18

15/16




Fig. 19




Fig. 20




Fig. 21

16/16

1 ATGCCAAGCGAACCGGCGATATACTAATTTCAACGCCCGTCTCGAAGGTTTCGTGGAAGACTGAACCTTCGACAGCCCGTA TILCSV CP (SEQ ID No 12)

1 ATGCCAAGCGAACCGGCGATATCTAATCTCTAATCCCGTCTGAAAGGTCCGTAAGGACTGAACCTTCGACAGCCCGTA TILCSV CP silencing minus (SEQ ID No 6)

81 TACCAGCCGTCCTGCTGCGCCCGCCCACTGTCCCAAGGCATCAAGCGCTCGATCATGGACCTTACAGGCCCATGTATCGAAGCCGCG TILCSV CP (SEQ ID No 12)

81 CACCTCTCTGTCGCGCTGCGCCCGCCCAAGTCCAGGGCACTAAGAGGCGCATCTTGGACATATACAGAGCCCTATGTACAGGAAACCCGA TILCSV CP silencing minus (SEQ ID No 6)

161 GGATGTACAGAAATGTPACAAAGCCCTGTGATACCTCCGGTTGTGAAGGTTCCCTGTAAAGTTCCTGTAAAGTTCGACGACGACGCT TILCSV CP (SEQ ID No 12)

161 GGATGTATAGGATGTATAGGATAGCCCTAGATGTGCTCTCAGGGTTGCGAAGGACCCCTGCAAGGTTGCAATCGTATGAGCAACGCT TILCSV CP silencing minus (SEQ ID No 6)

241 GATGACGTCGAAGCAATACCGGTTGTGTGCGTTGTCTGATGTAACTAGGGGTTCTCGTATTACTCATAGAGTTGGTAA TILCSV CP (SEQ ID No 12)

241 GACGATGTGAAGCAACCGGTTGTGTGCTGCTGTTGCTGATGTGACTAGACGGTTCAAGGTTCACTCAAGGGTGGAGAA TILCSV CP silencing minus (SEQ ID No 6)

321 ACGTTTTTGTATCAAGTCAATTTATATATTAGGAAGATTTGGATGGATGAAAACATATAAAACAATAATCATACTAACCC TILCSV CP (SEQ ID No 12)

321 CGCTTCTGTATAGTCAATTTATATATTGGGTAAGATCTGGATGGACGAACTATCAAGGAAACAGAAATCACTAATATCT TILCSV CP silencing minus (SEQ ID No 6)

401 AAGTGATGTTTTCTCTGTTTCGAGACCGAAGGCGCTTAAGGAACTAGTCCATGGATTTGGTCAAGTTTCTTAACTATGTT TILCSV CP (SEQ ID No 12)

401 AAGGTATGTTCTTCTGTTTCGAGAGTCGAAAGCCATAGGAAACGACCCATGGACATGGAGCTTGGCCAGGTTTAACTATGTTCT TILCSV CP silencing minus (SEQ ID No 6)

481 GATAATGAACCCAGTACTGCTACGGTGAAGAACGATTCAGGATAGGTATCAAGTAATGAGGAAGTTTCATGCTACCGGT TILCSV CP (SEQ ID No 12)

481 GATAACGAGCCATCTACTGCACTATGTGAAGAAAGGATTCGCTGATAGATATCAAGGATGATGAGGAAGTTTCATGCTACCGGT TILCSV CP silencing minus (SEQ ID No 6)

561 TGTGTGAGGTCCTCAGGGATGAAGGAGCAGTGTTCCTGAAGAGATTTTTTAAATAATAATACCCATGAGTTTATAATATC TILCSV CP (SEQ ID No 12)

561 GGTGTTGGTCTCTCTGGAATGAAGAGCACTGTCTGTGAAGATCTCTTTAAGATGCAACATCATGTCGTTATATAGCC TILCSV CP silencing minus (SEQ ID No 6)

641 ACCAAGAGCAGCGGAAGTATGAAATCATACTGAGATGCTTGTATTGTATATGGCTTGTACTCATGCTCTAACCCTA TILCSV CP (SEQ ID No 12)

641 ACCAGGAGCAAGCGAATATGAGAAATCACTGAAATCTTGTCTTATCATGGCTGTACCCAGCCATCTAATCTCCA TILCSV CP silencing minus (SEQ ID No 6)

721 GTGTACGCTACGTGTGAAAATACGTATTTATTTTATGATGCTGTAAACAAATTA TILCSV CP (SEQ ID No 12)

721 GTTTACGCAACGCTTAAGATCTCGTATCTATTTCTATGACGCTGTGACAAACCTAG TILCSV CP silencing minus (SEQ ID No 6)

Fig. 22

SEQUENCE LISTING

IAP20 Rec'd PCT/IT2004 17 NOV 2005

<110> ENEA-Ente per le Nuove Tecnologie e l'Ambiente
Consiglio Nazionale delle Ricerche

<120> Method for the preparation of transgenic plants characterised by
Geminivirus lasting resistance

<130> PCT25622

<140> RM2003A000242
<141> 2003-05-19

<150> RM2003A000242
<151> 2003-05-19

<160> 12

<170> PatentIn version 3.2

<210> 1
<211> 630
<212> DNA
<213> Geminivirus TYLCSV

<400> 1
atgccaagat caggctcggtt tagtatcaag gctaaaaatt atttccttac atatcccaa 60
tgtgatttaa caaaagaaaa tgcactttcc caaataacaa acctacaaac acccacaac 120
aaattattca tcaaaatttg cagagaacta catgaaaatg gggaacctca tctccatatt 180
ctcatccaat tcgaaggaaa atacaattgt accaatcaac gattcttcga cctgggatcc 240
ccaaccaggt cagcacattt ccatccgaac attcaggagg ctaaatacgag ctccgacgtc 300
aagtcctata tcgacaagga cggagatggt cttgaatggg gtactttcca gatcgacgga 360
cgatctgcta ggggaggaca acagacagcc aacgacgctt acgcaaaggc aattaacgca 420
ggaagtaagt cgcaggctct tgatgtaatt aaagaattag cgcctagaga ttacgttcta 480
cattttcata atataaatag taatttagat aaggttttcc aggtgcctcc ggcaccttat 540
gtttctcctt ttttatcttc ttctttcgat caagttcctg atgaacttga aactggggtt 600
tccgagaacg tcatggatgc cgctgcgcgg 630

<210> 2
<211> 630
<212> DNA
<213> Artificial

<220>
<223> TYLCSV Rep-210 modified sequence

<220>
<221> CDS
<222> (1)..(630)

<400> 2
atg cct aga tcc gga agg ttt agc atc aaa gct aag aat tac ttc ttg 48
Met Pro Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu
1 5 10 15
aca tac ccc aag tgt gac tta act aag gag aat gca ttg tcc cag ata 96
Pagina 1

Thr Tyr Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile
 20 25 30

act aac ttg caa act ccc act aac aag ttg ttc att aag att tgt agg 144
 Thr Asn Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg
 35 40 45

gaa ctt cat gag aat gga gaa cca cat ctt cat atc ttg ata cag ttc 192
 Glu Leu His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe
 50 55 60

gaa ggc aag tat aac tgc acc aac caa cgt ttc ttt gac ctt gtg tcc 240
 Glu Gly Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser
 65 70 75 80

cct acc aga tca gcc cat ttt cat cca aac atc cag ggt gct aag tcg 288
 Pro Thr Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser
 85 90 95

agt tca gac gtg aag tca tac att gac aaa gac ggc gat gtg ctc gag 336
 Ser Ser Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu
 100 105 110

tgg gga act ttt cag ata gac ggt cga tgc gct aga gga ggt cag caa 384
 Trp Gly Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln
 115 120 125

aca gct aac gat gca tac gct aag gct atc aac gct gga tcc aag tca 432
 Thr Ala Asn Asp Ala Tyr Ala Lys Ala Ile Asn Ala Gly Ser Lys Ser
 130 135 140

cag gca ctt gac gta atc aaa gag tta gct cct agg gat tat gtt ctt 480
 Gln Ala Leu Asp Val Ile Lys Glu Leu Ala Pro Arg Asp Tyr Val Leu
 145 150 155 160

cat ttc cat aac atc aac agc aat ttg gac aaa gtg ttc caa gtg cca 528
 His Phe His Asn Ile Asn Ser Asn Leu Asp Lys Val Phe Gln Val Pro
 165 170 175

ccg gct cct tac gtt tca cct ttc tta agt tct tca ttt gat cag gtt 576
 Pro Ala Pro Tyr Val Ser Pro Phe Leu Ser Ser Ser Phe Asp Gln Val
 180 185 190

cca gat gag ctt gag cat tgg gtg tcc gaa aac gtt atg gac gcc gca 624
 Pro Asp Glu Leu Glu His Trp Val Ser Glu Asn Val Met Asp Ala Ala
 195 200 205

gcg cgt
 Ala Arg
 210 630

<210> 3
 <211> 210
 <212> PRT
 <213> Artificial

<220>
 <223> Synthetic Construct

<400> 3

Met Pro Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu
 1 5 10 15

Thr Tyr Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile
 20 25 30

Thr Asn Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg
 35 40 45
 Glu Leu His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe
 50 55 60
 Glu Gly Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser
 65 70 75 80
 Pro Thr Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser
 85 90 95
 Ser Ser Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu
 100 105 110
 Trp Gly Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln
 115 120 125
 Thr Ala Asn Asp Ala Tyr Ala Lys Ala Ile Asn Ala Gly Ser Lys Ser
 130 135 140
 Gln Ala Leu Asp Val Ile Lys Glu Leu Ala Pro Arg Asp Tyr Val Leu
 145 150 155 160
 His Phe His Asn Ile Asn Ser Asn Leu Asp Lys Val Phe Gln Val Pro
 165 170 175
 Pro Ala Pro Tyr Val Ser Pro Phe Leu Ser Ser Ser Phe Asp Gln Val
 180 185 190
 Pro Asp Glu Leu Glu His Trp Val Ser Glu Asn Val Met Asp Ala Ala
 195 200 205
 Ala Arg
 210

<210> 4
 <211> 630
 <212> DNA
 <213> Artificial

<220>
 <223> TYLCSV Rep-210 modified sequence

<220>
 <221> CDS
 <222> (1)..(630)

<400> 4
 atg cct aga tcc gga agg ttt agc atc aaa gct aag aat tac ttc ttg
 Met Pro Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu
 1 5 10 15

48

aca tac ccc aag tgt gac tta act aag gag aat gca ttg tcc cag ata 96
 Thr Tyr Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile
 20 25 30
 act aac ttg caa act ccc act aac aag ttg ttc att aag att tgt agg 144
 Thr Asn Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg
 35 40 45
 gaa ctt cac gag aat gga gaa cca cat ctt cat atc ttg ata cag ttc 192
 Glu Leu His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe
 50 55 60
 gaa ggc aag tat aac tgc acc aac caa cgt ttc ttt gac ctt gtg tcc 240
 Glu Gly Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser
 65 70 75 80
 cct acc aga tca gcc cat ttt cat cca aac atc cag ggt gct aag tcg 288
 Pro Thr Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser
 85 90 95
 agt tca gac gtg aag tca tac att gac aaa gac ggg gat gtg ctc gag 336
 Ser Ser Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu
 100 105 110
 tgg gga act ttt cag ata gac ggt cga tgc gct aga gga ggt cag caa 384
 Trp Gly Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln
 115 120 125
 aca gca aac gat gca tac gct aag gct atc aac gct gga tcc aag tca 432
 Thr Ala Asn Asp Ala Tyr Ala Lys Ala Ile Asn Ala Gly Ser Lys Ser
 130 135 140
 cag gca ctt gac gta atc aaa gag tta gct cct agg gat tat gtt ctt 480
 Gln Ala Leu Asp Val Ile Lys Glu Leu Ala Pro Arg Asp Tyr Val Leu
 145 150 155 160
 cat ttc cat aac atc aac agc aat ttg gac aaa gtg ttc caa gtg cca 528
 His Phe His Asn Ile Asn Ser Asn Leu Asp Lys Val Phe Gln Val Pro
 165 170 175
 ccg gct cct tac gtt tca cct ttc tta agt tct tca ttt gat cag gtt 576
 Pro Ala Pro Tyr Val Ser Pro Phe Leu Ser Ser Ser Phe Asp Gln Val
 180 185 190
 cca gat gag ctt gag cat tgg gtg tct gaa aac gtt atg gac gcc gca 624
 Pro Asp Glu Leu Glu His Trp Val Ser Glu Asn Val Met Asp Ala Ala
 195 200 205
 gcc cgt
 Ala Arg
 210

<210> 5
 <211> 210
 <212> PRT
 <213> Artificial

<220>
 <223> Synthetic Construct

<400> 5

Met Pro Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu
 1 5 10 15

Thr Tyr Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile
 Pagina 4

20 25 30
 Thr Asn Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg
 35 40 45
 Glu Leu His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe
 50 55 60
 Glu Gly Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser
 65 70 75 80
 Pro Thr Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser
 85 90 95
 Ser Ser Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu
 100 105 110
 Trp Gly Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln
 115 120 125
 Thr Ala Asn Asp Ala Tyr Ala Lys Ala Ile Asn Ala Gly Ser Lys Ser
 130 135 140
 Gln Ala Leu Asp Val Ile Lys Glu Leu Ala Pro Arg Asp Tyr Val Leu
 145 150 155 160
 His Phe His Asn Ile Asn Ser Asn Leu Asp Lys Val Phe Gln Val Pro
 165 170 175
 Pro Ala Pro Tyr Val Ser Pro Phe Leu Ser Ser Ser Phe Asp Gln Val
 180 185 190
 Pro Asp Glu Leu Glu His Trp Val Ser Glu Asn Val Met Asp Ala Ala
 195 200 205
 Ala Arg
 210

<210> 6
 <211> 774
 <212> DNA
 <213> Artificial

<220>
 <223> TYLCSV Coat Protein modified sequence

<220>
 <221> CDS
 <222> (1)..(774)

<400> 6
 atg cca aag aga act ggt gat att cta atc tca act ccc gtg tct aag
 Met Pro Lys Arg Thr Gly Asp Ile Leu Ile Ser Thr Pro Val Ser Lys
 1 5 10 15

gtg cgt agg aga ctt aac ttt gac tct ccg tac acc tct cgt gca gct Val Arg Arg Arg Leu Asn Phe Asp Ser Pro Tyr Thr Ser Arg Ala Ala 20 25 30	96
gct ccc aca gtc cag ggc att aag agg cga tct tgg aca tac aga cct Ala Pro Thr Val Gln Gly Ile Lys Arg Arg Ser Trp Thr Tyr Arg Pro 35 40 45	144
atg tac agg aaa ccg agg atg tat agg atg tat cgt agc cca gat gtg Met Tyr Arg Lys Pro Arg Met Tyr Arg Met Tyr Arg Ser Pro Asp Val 50 55 60	192
cct cct ggt tgc gaa gga ccc tgc aag gtg caa tcg tat gag caa cgt Pro Pro Gly Cys Glu Gly Pro Cys Lys Val Gln Ser Tyr Glu Gln Arg 65 70 75 80	240
gac gat gtg aag cac acc gga gtt gtt cgt tgc gtt tct gat gtg act Asp Asp Val Lys His Thr Gly Val Val Arg Cys Val Ser Asp Val Thr 85 90 95	288
aga ggt tca ggt atc act cac agg gtg gga aag cgt ttc tgt att aag Arg Gly Ser Gly Ile Thr His Arg Val Gly Lys Arg Phe Cys Ile Lys 100 105 110	336
tct att tac ata ttg ggt aag atc tgg atg gac gag aat atc aag aaa Ser Ile Tyr Ile Leu Gly Lys Ile Trp Met Asp Glu Asn Ile Lys Lys 115 120 125	384
cag aat cac act aat cag gtt atg ttc ttt ctt gtg cga gat cga aga Gln Asn His Thr Asn Gln Val Met Phe Phe Leu Val Arg Asp Arg Arg 130 135 140	432
cca tac gga acc agc cca atg gac ttc ggc cag gtg ttt aat atg ttc Pro Tyr Gly Thr Ser Pro Met Asp Phe Gly Gln Val Phe Asn Met Phe 145 150 155 160	480
gat aac gag cca tct act gca act gtg aaa aat gat ttg cgt gat aga Asp Asn Glu Pro Ser Thr Ala Thr Val Lys Asn Asp Leu Arg Asp Arg 165 170 175	528
tat cag gtg atg aga aag ttc cat gca acg gtg gtt ggt ggt cct tct Tyr Gln Val Met Arg Lys Phe His Ala Thr Val Val Gly Gly Pro Ser 180 185 190	576
gga atg aaa gag caa tgt ctt ctg aaa aga ttc ttt aag atc aac act Gly Met Lys Glu Gln Cys Leu Leu Lys Arg Phe Phe Lys Ile Asn Thr 195 200 205	624
cat gtc gtc tat aac cac cag gag caa gcg aaa tat gag aat cac act His Val Val Tyr Asn His Gln Glu Gln Ala Lys Tyr Glu Asn His Thr 210 215 220	672
gaa aat gct ttg ttg tta tac atg gcc tgt acc cac gca tct aat cca Glu Asn Ala Leu Leu Leu Tyr Met Ala Cys Thr His Ala Ser Asn Pro 225 230 235 240	720
gtt tac gca acg ctt aag atc cgt atc tat ttc tat gac gct gtg aca Val Tyr Ala Thr Leu Lys Ile Arg Ile Tyr Phe Tyr Asp Ala Val Thr 245 250 255	768
aac tag Asn	774

<210> 7
<211> 257

<212> PRT
 <213> Artificial

<220>
 <223> Synthetic Construct

<400> 7

Met Pro Lys Arg Thr Gly Asp Ile Leu Ile Ser Thr Pro Val Ser Lys
 1 5 10 15

Val Arg Arg Arg Leu Asn Phe Asp Ser Pro Tyr Thr Ser Arg Ala Ala
 20 25 30

Ala Pro Thr Val Gln Gly Ile Lys Arg Arg Ser Trp Thr Tyr Arg Pro
 35 40 45

Met Tyr Arg Lys Pro Arg Met Tyr Arg Met Tyr Arg Ser Pro Asp Val
 50 55 60

Pro Pro Gly Cys Glu Gly Pro Cys Lys Val Gln Ser Tyr Glu Gln Arg
 65 70 75 80

Asp Asp Val Lys His Thr Gly Val Val Arg Cys Val Ser Asp Val Thr
 85 90 95

Arg Gly Ser Gly Ile Thr His Arg Val Gly Lys Arg Phe Cys Ile Lys
 100 105 110

Ser Ile Tyr Ile Leu Gly Lys Ile Trp Met Asp Glu Asn Ile Lys Lys
 115 120 125

Gln Asn His Thr Asn Gln Val Met Phe Phe Leu Val Arg Asp Arg Arg
 130 135 140

Pro Tyr Gly Thr Ser Pro Met Asp Phe Gly Gln Val Phe Asn Met Phe
 145 150 155 160

Asp Asn Glu Pro Ser Thr Ala Thr Val Lys Asn Asp Leu Arg Asp Arg
 165 170 175

Tyr Gln Val Met Arg Lys Phe His Ala Thr Val Val Gly Gly Pro Ser
 180 185 190

Gly Met Lys Glu Gln Cys Leu Leu Lys Arg Phe Phe Lys Ile Asn Thr
 195 200 205

His Val Val Tyr Asn His Gln Glu Gln Ala Lys Tyr Glu Asn His Thr
 210 215 220

Glu Asn Ala Leu Leu Leu Tyr Met Ala Cys Thr His Ala Ser Asn Pro
 225 230 235 240

Val Tyr Ala Thr Leu Lys Ile Arg Ile Tyr Phe Tyr Asp Ala Val Thr
245 250 255

Asn

<210>	8
<211>	447
<212>	DNA
<213>	Artificial

<220>
<223> TYLCSV Rep 130 sequence

<220>
<221> CDS
<222> (51) .. (443)

```

<220>
<221> misc_feature
<222> (231)..(231)
<223> Point mutation from C (Rep-210 wild-type) to T

```

```
<220>
<221> misc_feature
<222> (233)..(233)
<223> Point mutation from C (Rep 210 wild-type) to G
```

<400> 8
ggatccccct ggatactttg agtgtcccc gattcagaac gacagcaaaa atg cca 56
Met Pro
1

aga tca ggt cgt ttt agt atc aag gct aaa aat tat ttc ctt aca tat 104
Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu Thr Tyr
5 10 15

ccc aaa tgt gat tta aca aaa gaa aat gca ctt tcc caa ata aca aac 152
Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile Thr Asn
20 25 30

cta caa aca ccc aca aac aaa tta ttc atc aaa att tgc aga gaa cta 200
Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg Glu Leu
35 40 45 50

cat gaa aat ggg gaa cct cat ctc cat att ttg atc caa ttc gaa gga 248
His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe Glu Gly
55 60 65

aaa tac aat tgt acc aat caa cga ttc ttc gac ctg gta tcc cca acc 296
Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser Pro Thr
70 75 80

agg tca gca cat ttc cat ccg aac att cag gga gct aaa tcg agc tcc 344
Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser Ser Ser
85 90 95

gac gtc aag tcc tat atc gac aag gac gga gat gtt ctt gaa tgg ggt 392
Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu Trp Gly
100 105 110

act ttc cag atc gac gga cga tct gct agg gga gga caa cag aca gcc 440
Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln Thr Ala
115 120 125 130

tga attc

447

<210> 9
 <211> 130
 <212> PRT
 <213> Artificial

<220>
 <223> Synthetic Construct

<400> 9

Met Pro Arg Ser Gly Arg Phe Ser Ile Lys Ala Lys Asn Tyr Phe Leu
 1 5 10 15

Thr Tyr Pro Lys Cys Asp Leu Thr Lys Glu Asn Ala Leu Ser Gln Ile
 20 25 30

Thr Asn Leu Gln Thr Pro Thr Asn Lys Leu Phe Ile Lys Ile Cys Arg
 35 40 45

Glu Leu His Glu Asn Gly Glu Pro His Leu His Ile Leu Ile Gln Phe
 50 55 60

Glu Gly Lys Tyr Asn Cys Thr Asn Gln Arg Phe Phe Asp Leu Val Ser
 65 70 75 80

Pro Thr Arg Ser Ala His Phe His Pro Asn Ile Gln Gly Ala Lys Ser
 85 90 95

Ser Ser Asp Val Lys Ser Tyr Ile Asp Lys Asp Gly Asp Val Leu Glu
 100 105 110

Trp Gly Thr Phe Gln Ile Asp Gly Arg Ser Ala Arg Gly Gly Gln Gln
 115 120 125

Thr Ala
 130

<210> 10
 <211> 30
 <212> DNA
 <213> Artificial

<220>
 <223> Primer

<220>
 <221> misc_feature
 <222> (1)..(30)
 <223> Primer for PCR C4 mutagenesis

<400> 10
 ctcacctcca tttttgatc caattcgaag

30

<210> 11

<211> 30
 <212> DNA
 <213> Artificial

<220>
 <223> Primer

<220>
 <221> misc_feature
 <222> (1)..(30)
 <223> Primer for PCR C4 mutagenesis

<400> 11
 cttcgaattg gatcaaaata tggagatgag 30

<210> 12
 <211> 774
 <212> DNA
 <213> Geminivirus TYLCSV

<400> 12
 atgccgaagc gaaccggcga tataactaatt tcaacgcccg tctcgaagggt tcgtcgaaga 60
 ctgaacttcg acagcccgtg taccagccgt gctgctgccc cactgtcca aggcacaaag 120
 cgtcgatcat ggacttacag gcccatgtat cgaaagccgc ggatgtacag aatgtacaga 180
 agccctgatg tacctccggg ttgtgaagggt ccctgtaaag tgcagtcgta cgagcagcgt 240
 gatgacgtca agcataaccg tgtgtgctgt tgtgttagtg atgtaactag gggttctggt 300
 attactcata gagttggtaa acgtttttgt atcaagtcaa tttatatatt aggaaagatt 360
 tggatggatg aaaacataaa aaaacaaaat cataactaacc aagtgatggt tttccttggt 420
 cgagaccgaa ggccttatgg aactagtcct atggattttg gtcaagtttt taacatgttt 480
 gataatgaac ccagtactgc tacgggtgaag aacgacttac gggataggta tcaagtaatg 540
 aggaagtttc atgctacggt tgttggagggt ccgtcaggga tgaaggagca gtgtttgctg 600
 aagagatttt ttaaaattaa tacccatgta gtttataatc accaagagca ggcgaagtat 660
 gaaaatcata ctgagaatgc cttgttattg tatatggctt gtactcatgc ttctaaccga 720
 gtgtacgcta cgttgaaaat acgtatttat ttttatgatg ctgtaacaaa ttaa 774

INTERNATIONAL SEARCH REPORT

International Application No
PCT/AT2004/000287

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/82 C07K14/01

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, Sequence Search, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/43520 A (ABOUZID AHMED M ; HIEBERT ERNEST (US); HUNTER WAYNE B (US); POLSTON JA) 27 July 2000 (2000-07-27) the whole document	1-43
A	BRUNETTI ANGELA ET AL: "Transgenically expressed T-Rep of tomato yellow leaf curl sardinia virus acts as a trans-dominant-negative mutant, inhibiting viral transcription and replication" JOURNAL OF VIROLOGY, vol. 75, no. 22, November 2001 (2001-11), pages 10573-10581, XP002300782 ISSN: 0022-538X the whole document	1-43

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

15 October 2004

Date of mailing of the international search report

02/11/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mundel, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US2004/000287

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>NORIS E ET AL: "Resistance to tomato yellow leaf curl geminivirus in Nicotiana benthamiana plants transformed with a truncated viral C1 gene"</p> <p>VIROLOGY, ACADEMIC PRESS, ORLANDO, US, vol. 224, no. 1, 1996, pages 130-138, XP002142359</p> <p>ISSN: 0042-6822</p> <p>the whole document</p> <p>-----</p>	1-43

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter. Appl. No.
PCT/JP2004/000287

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0043520 A	27-07-2000	AU 2856400 A WO 0043520 A2	07-08-2000 27-07-2000
<hr/>			